

Digital interventions to reduce social isolation and loneliness in older adults: An evidence and gap map

Review information

Authors

Vivian Welch, Elizabeth Ghogomu, Victoria Barbeau, Sierra Dowling, Rebecca Doyle, Ella Beveridge, Elisabeth Boulton, Payaam Desai, Jimmy Huang, Nour Elmostekawy, Tarannum Hussain, Arpana Wadhvani, Sabrina Boutin, Niobe Haitas, Dylan Kneale, Douglas M. Salzwedel, Roger Simard, Paul Hebert, Christopher Mikton

Contact person

Vivian Welch

Methods Centre, Bruyère Research Institute,
Ottawa, Canada.

Email: vwelch@campbellcollaboration.org

Abstract

Background

Social isolation and loneliness are more common in older adults and are associated with a serious impact on their well-being, mental health, physical health, and longevity. They are a public health concern highlighted by the COVID-19 pandemic restrictions, hence the need for digital technology tools to enable remotely delivered interventions to alleviate the impact of social isolation and loneliness during the COVID-19 restrictions.

Objectives

To map available evidence on the effects of digital interventions to mitigate social isolation and/or loneliness in older adults in all settings except hospital settings.

Search methods

We searched the following databases from inception to 16 May 2021 with no language restrictions: Ovid MEDLINE, Embase, APA PsycInfo via Ovid, CINAHL via EBSCO, Web of Science via Clarivate, ProQuest (all databases), International Bibliography of the Social Sciences (IBSS) via ProQuest, EBSCO (all databases except CINAHL), Global Index Medicus, and Epistemonikos.

Selection criteria

Titles and abstracts and full text of potentially eligible articles were independently screened in duplicate following the eligibility criteria.

Data collection and analysis

We developed and pilot tested a data extraction code set in Eppi-Reviewer and data was individually extracted and coded.

Main results

We included 200 articles (103 primary studies and 97 systematic reviews) that assessed the effects of digital interventions to reduce social isolation and/or loneliness in older adults. Most of the systematic reviews (72%) were classified as critically low quality, only 2% as high quality and 25% were published since the COVID-19 pandemic.

The evidence is unevenly distributed with clusters predominantly in high-income countries and none in low-income countries. The most common interventions identified are digital interventions to enhance social interactions with family and friends and the community via videoconferencing and telephone calls. Digital interventions to enhance social support, particularly socially assistive robots, and virtual pets were also common. Most interventions focused on reducing loneliness and depression and improving quality of life of older adults.

Major gaps were identified in community level outcomes and process indicators. No included studies or reviews assessed affordability or digital divide although the value of accessibility and barriers caused by digital divide were discussed in three primary studies and three reviews. Adverse effects were reported in only two studies and six reviews.

No study or review included participants from the LGBTQIA2S+ community and only one study restricted participants to 80 years and older. Very few described how at-risk populations were recruited or conducted any equity analysis to assess differences in effects for populations experiencing inequities across PROGRESS-Plus categories.

Authors' conclusions

The restrictions placed on people during the pandemic have shone a spotlight onto social isolation and loneliness, particularly for older adults. This evidence and gap map shows available evidence on the effectiveness of digital interventions for reducing social isolation or loneliness in older adults.

Although the evidence is relatively large and recent, it is unevenly distributed and there is need for more high-quality research. This map can guide researchers and funders to consider areas of major gaps as priorities for further research.

Abbreviations and acronyms

EGM: evidence and gap map

LGBTQIA2S+: lesbian, gay, bisexual, transgender, queer (or sometimes questioning), intersex, asexual, and two-spirited

PICO: population, intervention, comparison, outcome

PROGRESS-Plus: Place of residence (urban/rural), Race/ethnicity/culture and language, Occupation, Gender or sex, Religion, Occupation, Socioeconomic status, Social capital and Plus factors

WHO: World Health Organization

Plain language summary

The evidence for digital interventions to reduce social isolation and loneliness in older adults is unevenly distributed geographically and by outcomes, and most of the reviews are of critically low quality.

Mapping the evidence for digital interventions to reduce social isolation and loneliness in older adults shows that many of the systematic reviews are of critically low quality and most come from high-income countries with sparse reporting of community level outcomes.

What is this EGM about?

Social isolation and loneliness are common in older adults and have been highlighted during the COVID-19 pandemic. They pose a significant concern because of serious impact on older adults' well-being, mental health, physical health, and longevity.

Older adults can use digital interventions to maintain existing connections or develop new connections especially during the COVID-19 pandemic with social distancing and lockdown measures in place. Decision makers need to know which digital interventions can reduce social isolation and loneliness in older adults.

What is the aim of this EGM?

To show available evidence from systematic reviews and primary studies on the effects of digital interventions to reduce social isolation and/or loneliness among older adults in all settings except hospital settings.

What studies are included?

The EGM includes 200 articles (97 systematic reviews and 103 primary studies) that assessed how digital interventions can reduce social isolation and loneliness in older adults. The studies had to report the effect of digital interventions and could come from any region.

What are the main findings of this gap map?

The evidence is unevenly distributed geographically, with most from high-income countries and none from low-income countries. Over 70% of the systematic reviews have critically low quality and 25% have been published since the pandemic began.

The most common interventions are digital interventions to enhance social interactions with family, friends and the community via videoconferencing and telephone calls. Digital interventions to enhance social support, particularly socially assistive robots, and virtual pets were also common. Most interventions focus on reducing loneliness and depression and improving quality of life of older adults.

Community level outcomes and process indicators are hardly reported, and no included studies or reviews assess affordability or lack of accessibility, although the value of accessibility and barriers caused by lack of accessibility were discussed in three studies and three reviews.

Adverse effects are reported in very few studies and reviews.

Participants from the LGBTQIA2S+ community are not included in any study or review and only one study restricted participants to 80 years and older. Very few studies or reviews describe how at-risk populations were recruited or conduct any equity analysis to assess differential effects for populations experiencing inequities across PROGRESS-Plus factors.

What do the findings of the map mean?

This evidence and gap map is a first step towards identifying which digital interventions are effective for reducing social isolation and loneliness in older adults and contributes to the longer-term aim of building an evidence architecture for the field, whereby the use of evidence becomes institutionalized in policy and practice.

Although the evidence is relatively large and recent, it is unevenly distributed and there is need for more high-quality research. This map can guide researchers and funders to consider areas of major gaps as priorities for further research.

How up-to-date is this map?

The authors searched for studies published up to May 2021.

Background

Introduction

The problem, condition or issue

A large body of research shows that social isolation and loneliness are associated with a serious impact on older adults' well-being, mental health, physical health, and longevity (Leigh-Hunt et al., 2017; Menec et al., 2020). Their effect on mortality is comparable to, or even greater than, other well-established risk factors such as smoking, obesity, and physical inactivity (Holt-Lunstad et al., 2015; Ibarra et al., 2020; Menec et al., 2020; Windle et al., 2019).

Social isolation and loneliness are more common in older adults and are described as multidimensional concepts with different methods of measurement leading to variations in the prevalence. The prevalence ranges from 5% to 43% depending on the study and region (Chen & Schulz, 2016; Donovan & Blazer, 2020; Ibarra et al., 2020; Leigh-Hunt et al., 2017). Risk factors include living alone, impaired mobility, experiencing a major life transition change (e.g., loss of spouse or other primary network members), limited income or resources, cognitive impairment, inadequate social support, and geographic location (Cohen-Mansfield & Perach, 2015; Donovan & Blazer, 2020; Findlay, 2003; Ibarra et al., 2020).

Although they are related, social isolation and loneliness are two distinct concepts, and one may occur without the other. Social isolation is the objective state of lack of interactions with others and the wider community, or lack of social relationships (Donovan & Blazer, 2020; Ibarra et al., 2020; Leigh-Hunt et al., 2017; Menec et al., 2020). Loneliness is the subjective painful feeling of the absence of a [social network](#) or a companion, or perception of unmet emotional and social

needs resulting from a mismatch between the desired and actual experience of the quality or quantity of social relationships (Cacioppo et al., 2009, 2014; Menec et al., 2020; Perlman & Peplau, 1981; Prohaska et al., 2020; World Health Organization, 2021). Therefore, an individual can have a social network and be lonely, or a socially isolated individual may not feel lonely. An understanding of the differences in these concepts is important for research in the development of appropriate and effective interventions, and standardizing outcome measurements and also to guide the choice of appropriate interventions for socially isolated or lonely individuals (Fakoya et al., 2020; World Health Organization, 2021).

Social isolation and loneliness among older adults are priority public health problems, as well as national and international policy issues, due to the negative impact on their mental and physical health and longevity (Cattan et al., 2005; Gardiner et al., 2018; Shah et al., 2020, 2021; World Health Organization, 2020, 2021). The World Health Organization (WHO) decided, as part of the Decade of Healthy Ageing, to address social isolation and loneliness as a priority issue that cuts across the main action areas of the Decade (World Health Organization, 2020). It is also increasingly being recognized as a public health concern due to the social distancing measures during the COVID-19 pandemic (Brooke & Jackson, 2020; Shah et al., 2020, 2021; Williams et al., 2021). For example, the average person's daily number of contacts was reduced by up to 74% and almost one quarter of adults in the UK experienced loneliness when living under lockdown (Williams et al., 2021). Hence the need for digital technology tools to enable remotely delivered interventions to alleviate the impact of social isolation and loneliness during the COVID-19 restrictions.

There are challenges associated with access to digital interventions and the use of remotely delivered interventions to reduce social isolation and loneliness. Disparities in access to digital interventions and the use of remotely delivered interventions is a growing concern, especially for older adults and during the COVID-19 restrictions (Budd et al., 2020; Jopling, 2020; Watts, 2020; Williams et al., 2021). Many older adults lack digital skills and the confidence to access online services and support. Other barriers are affordability and accessibility of technology, broadband or Wi-Fi, data poverty (i.e., lack of accessibility to wireless internet connection), geographic divide (rural and urban, high income and low- and middle-income countries). Concerns with digital technology use have also been raised regarding privacy invasion, legal, ethical and clinical data governance through data sharing and access to information (Budd et al., 2020; NASEM 2020). Ethical concerns with any intervention to prevent social isolation and loneliness include issues with accessibility, acceptability, cost, feasibility, autonomy, and

informed concern. Privacy concerns may intersect with accessibility issues due to cognitive impairment or involvement of other family members or caregivers. Informed consent provided at the beginning of research may not cover the future use of data created, raising issues with ongoing consent and the ability to withdraw consent. There is also the risk of digital technology exacerbating rather than reducing social isolation, for example, social robots replacing meaningful human contact can result in increased sense of loneliness, deception and infantilization of older adults (NASEM, 2020). Equitable access and support are key in addressing the digital divide.

The intervention

A wide variety of interventions have been developed to reduce social isolation or loneliness among older adults. These interventions use different strategies and target different aspects such as facilitating social connections or service provision. They are implemented at different levels such as one-on-one or group focused. Although several systematic reviews have evaluated the effectiveness of different types of interventions for social isolation and loneliness in older adults, their findings have sometimes been conflicting (Cattan et al., 2005; Cohen-Mansfield & Perach, 2015; Dickens et al., 2011; Findlay, 2003; Gardiner et al., 2018; Hagan et al., 2014; C. Victor et al., 2018).

Digital interventions have become a particular focus of interest, due partly to the social distancing and lockdown measures introduced to combat the COVID-19 pandemic, and to the rapidly increasing role technology – particularly the internet, mobile devices, social media and Internet of things (IoT) – has played in the last 10-15 years in mediating social relations (Boulton et al., 2020; Brooke & Jackson, 2020; Budd et al., 2020; Falk, 2021; Shah et al., 2020; UCLG, 2020; World Health Organization, 2021; Zanella et al., 2020). They have been used in different sectors (e.g., health care, social services, the community) and in various ways, including digital epidemiological surveillance, rapid case identification, interruption of community transmission, public communication, and provision of clinical care and income support and livelihood opportunities in the COVID-19 crisis.

Digital interventions have also been used to mitigate social isolation or loneliness in older adults, by facilitating social interaction or by delivering programs or services (Boulton et al., 2020; Chen & Schulz, 2016; Chipps et al., 2017; Findlay, 2003; Ibarra et al., 2020; Khosravi et al., 2016; Noone et al., 2020; Shah et al., 2021; Thangavel et al., 2022). They have generally been described as technology-based interventions to improve communication and social

connection among older adults and there is no clear framework for their categorization (Fakoya et al., 2020). For example, they have been categorized as one-on-one or group-based interventions (Cohen-Mansfield & Perach, 2015; Dickens et al., 2011; Masi et al., 2011; Poscia et al., 2018) or based on four strategies or type (Masi et al., 2011) as:

- [interventions for improving social skills](#) (e.g., computer and internet training and use with a focus on reducing social isolation or loneliness; online university of the third age);
- [interventions for enhancing social support](#) that offer regular contacts, care, or companionship (e.g., telecare with a component to improve social connections, personal reminder information and social management systems (PRISMS), social robots or virtual pets, video games, 3D virtual environments or virtual spaces with trained coaches, conversational agents, or messaging capabilities);
- [interventions for enhancing social interaction](#) (videoconferencing, supported video communication, internet chat facilities, social networking sites, online discussion groups and forums, telephone befriending);
- [interventions for social cognitive training](#) (low intensity psychosocial interventions, internet-delivered cognitive behavioral therapy (CBT), mindfulness interventions).

In mapping the body of available evidence, we categorized interventions by strategies to enable exclusive coding of interventions in categories and subcategories such that an intervention will fit into a single subcategory and not overlap with another on the evidence and gap map.

Why it is important to develop the EGM

Several recent reviews of digital interventions for reducing social isolation and loneliness among older adults indicate there is growing research in this topic area most likely due to the ageing population (Boulton et al., 2020; Chen & Schulz, 2016; Chipps et al., 2017; Findlay, 2003; Ibarra et al., 2020; Khosravi et al., 2016; Noone et al., 2020; Shah et al., 2021). In addition, the COVID-19 pandemic restrictions have led to a dramatic expansion in the demand for digital technology interventions by people without access, including older adults, for the provision of basic services like healthcare, education, and connections with other people (UCLG, 2020).

Although there is a very wide range of such interventions, findings on their effectiveness, have sometimes been inconsistent (World Health Organization, 2021). The body of evidence supporting their use is rapidly expanding, dispersed and uneven with lack of consistent terminology. Therefore, the best use of resources at this point for building the [evidence](#)

[architecture](#) needed would be to develop an evidence and gap map on digital interventions to reduce social isolation and loneliness among older adults. This evidence and gap map will collate the evidence and display clusters of evidence and gaps in evidence that will serve as a resource to guide prioritization of further research and increase the accessibility and use of evidence for informed decision making by stakeholders including citizens, patients, caregivers, health and social care providers, policy makers and researchers.

Existing evidence and gap maps and/or relevant systematic reviews

Recent reviews of digital interventions suggest that a) there is a very wide range of such interventions; b) findings on their effectiveness, although sometimes positive, are frequently mixed, inconclusive or uncertain; and c) the technologies involved are developing rapidly (e.g. artificial intelligence, conversational agents, 3D virtual environments, video-games, social networking tools) (Boulton et al., 2020; Chen & Schulz, 2016; Chipps et al., 2017; Ibarra et al., 2020; Khosravi et al., 2016; Noone et al., 2020; Shah et al., 2021).

There is an evidence and gap map on specific remotely delivered interventions (i.e., befriending, social support, and low intensity psychosocial interventions) to reduce social isolation and loneliness among older adults (Boulton et al., 2020). It is based on a rapid review of reviews with systematic review evidence on befriending, social support, and low intensity psychosocial interventions that are delivered remotely to older adults, excluding caregivers. Study-level evidence is limited to 18 individual studies in the five included systematic reviews.

Our evidence and gap map will be more comprehensive with a broader scope of all types of digital interventions for older adults including older caregivers. It will examine up to date evidence from systematic reviews as well as primary studies and map available evidence to identify gaps and clusters in interventions and outcomes assessed.

Objectives

The aim is to map available evidence on the effects of digital interventions to mitigate social isolation and/or loneliness in older adults in all settings except hospital settings.

Methods

We followed the Campbell Collaboration guidance for producing an evidence and gap map (White et al., 2020) described in the evidence and gap map protocol for this project (Welch et al., 2022).

Evidence and gap map: definition and purpose

Evidence and gap maps are a systematic evidence synthesis product with a visual presentation of existing evidence relevant to a specific research question (Snilstveit et al., 2013; White et al., 2020). They display areas with collections or gaps in evidence and the quality of available evidence.

The evidence and gap map is typically a two dimensional matrix with interventions as row headings and outcomes as column headings (Snilstveit et al., 2016; White et al., 2020). Each cell within the matrix shows the studies with evidence on the corresponding intervention and outcome. This map identifies areas of evidence as well as any gaps in research related to using digital interventions for social isolation and/or loneliness among older adults.

Framework development and scope


We developed an intervention-outcome framework for this evidence and gap map through a consultative process with stakeholders and adaptation of existing frameworks from systematic reviews, conceptual papers, and reports from stakeholder organizations.

A refined version of the World Health Organization (WHO) Classification of Digital Health Interventions framework (World Health Organization, 2018) was initially considered at the Stakeholder consultation meeting on April 8, 2021. The WHO framework was developed to categorize the different ways in which digital and mobile technologies are used to support healthcare. The stakeholders found the typology of interventions to be too healthcare focused. The consensus was that a more user intuitive typology of interventions was needed to ensure the useability of this evidence and gap map for a larger audience including older adults. A needs-based approach was preferred as interventions are most effective when they meet the needs and specific circumstances of the older adults (Abdi et al., 2019; Findlay, 2003; ten Bruggencate et al., 2019; World Health Organization, 2020).

We identified other relevant frameworks from existing reviews and conceptual papers. We chose two frameworks which used a needs-based approach (Jopling, 2020) and a strategy-based approach (Masi et al., 2011) to address social isolation and loneliness and adapted them for our evidence and gap map.

The needs-based framework (Jopling, 2020) considers approaches to address loneliness and social isolation that are used in communities to achieve three outcomes: maintain and improve existing relationships or connections, support people to develop new connections, and to change negative thinking about their relationships. The approaches include connector services that reach out to understand the needs of older adults and provide support to meet the needs, gateway infrastructures through which people can connect with others, direct solutions or interventions to reduce loneliness and social isolation, and system-level approaches that create environments in communities to facilitate tackling loneliness and social isolation (Figure 1).

Figure 1: Needs-based approach framework (Adapted from Jopling et al. 2020)



The intervention categories in this framework do not provide mutually exclusive categorization of digital interventions. For example, many digital interventions such as computer and internet training, video chats, online cognitive behavioral therapy may be one-to-one, or group based. Hence the need for the second framework.

The strategy-based model (Masi et al., 2011) describes strategies used in loneliness reduction interventions based on the understanding of the nature of loneliness and social isolation and how they affect people (Figure 2). Interventions were also categorized based on the format or level of delivery (as one-on-one or group interventions) or mode of delivery (technology-based and non-technology-based interventions).

Figure 2: Strategy-based approach framework (Adapted from Masi et al. 2011)

Intervention format	Intervention mode	Intervention strategy (type)
<ul style="list-style-type: none"> • Individual-based (implemented on a one-on-one basis) • Group-based (more than one person participated) 	<ul style="list-style-type: none"> • Technology-based (e.g., telephone, computer used to facilitate the intervention) • Non-technology-based 	<ul style="list-style-type: none"> • Social skills training (if the intervention focused on improving participants' interpersonal communication skills) • Enhanced social support (if the intervention offered regular contacts, care, or companionship) • Social access (if the intervention increased opportunity for participants to engage in social interaction) • Social cognitive training (if the intervention focused on changing participants' social cognition)

We used an intervention-outcome framework where digital interventions of interest were coded by the strategies to reduce loneliness and social isolation: strategies for 1) improving social skills, 2) enhancing social support, 3) enhancing social interaction, 4) social cognitive training, and 5) multicomponent strategies; as well as by the type of intervention (e.g., computer and internet training to reduce social isolation and loneliness, video chats, telephone befriending, telecare with a component to improve social connections, online cognitive behavioral therapy). See [Appendix 1](#) for a glossary of key concepts.

Since the framework was bi-dimensional (interventions and outcomes), the needs of socially isolated and lonely older adults were used as a filter on the map and coded interventions were mapped to the needs.

Outcomes

The impacts of interventions to prevent social isolation and loneliness have been measured at different levels – individual, community or societal, and process and implementation (Windle et al., 2012). In our framework, we considered outcomes that have been identified as indicators of

[social connection](#) and they were categorized based on the impact and level of influence of the interventions:

- individual outcomes – loneliness, social isolation, social connectedness, quality of life, anxiety/depression, confidence level, information, communication and technologies (ICT) knowledge and experience, adverse effects;
- community outcomes – [social support](#), [social engagement](#), [social cohesion](#), [social capital](#), [digital divide](#) (disparities in access to technological interventions);
- process indicators – acceptance, adherence, technology use, feasibility, cost-effectiveness, affordability, barriers.

Stakeholder engagement

We convened an advisory board of 30 stakeholders from organizations such as the International Red Cross, Canadian Red Cross, Agewell, Canadian Frailty Network, HelpAge, CanAge, Centre for Ageing Better, United Nation Department of Social and Economic Affairs, United Nations Fund for Population Activities (UNFPA), and the World Health Organization (WHO). The group of stakeholders included representatives of these key organizations, policymakers, and academics with an interest in mitigating social isolation and loneliness in older adults. The advisory board provided comments on the intervention-outcome framework. The WHO Classification of Digital Health Interventions framework was considered. Stakeholders suggested a simplified framework to fit the purpose of this evidence and gap map. The framework was revised based on their feedback, and stakeholders were consulted by email for their feedback on the revised framework included in this review.


We consulted with four citizens in two citizen focus groups between June and August 2021. Some iterations were suggested, i.e., coding for interventions related to the need of finding purpose in later life, and capturing interventions related to recreation and physical activity. Affordability and access to technology were recommended for consideration as outcomes in the framework.

We created an anonymous survey with four questions and invited stakeholders including citizens by email to respond to the survey questions and provide their feedback on the revised framework and draft map.

Conceptual framework

Our conceptual framework (Figure 3) was based on the understanding of the needs of older adults, how social isolation and loneliness can occur and how they affect older adults' well-being. The relationship between these variables can be explained by a potential pathway of effect illustrated in the conceptual framework going from risk factors and needs assessment for older adults to interventions, the mechanisms of change, and process indicators and outcomes.

Figure 3: Conceptual framework



Population targeted by interventions

Older adults are more susceptible to experiencing social isolation and loneliness, but risk exposure may vary with individual contexts. Older adults at risk of isolation and loneliness have been identified by their age, gender, place of residence or other factors (Elder & Retrum, 2012; Fakoya et al., 2020; NASEM 2020). Interventions may target all older adults regardless of their risk (universal) or a subpopulation of older adults who are at risk e.g., those living in nursing homes (selective) or may target those who are socially isolated or lonely (indicated).

Risk factors for social isolation and loneliness

Social isolation and loneliness have been associated with low social support and a disruption in social interactions established with other people at any level (individual, group, community, and

societal or system) which can lead to unmet needs (Abdi et al., 2019; Donovan & Blazer, 2020; Tomaka et al., 2006). Major changes in life such as change or loss of social network, social participation or role, physical health, mental health and financial resources can also lead to social isolation and loneliness (Donovan & Blazer, 2020; Newall & Menec, 2019; Victor et al., 2005). Other risk factors for social isolation and loneliness include living alone, societal factors (racism, language barriers, ageism, social distancing and restrictions) and the physical environment (inaccessible location or community setting) (Berkman et al., 2000; DeGood, 2011; Donovan & Blazer, 2020).

Needs assessment for older adults

Ageing is associated with a decline in physical and cognitive health, difficulty with mobility, activities of daily living and household routines which put older adults at risk of experiencing needs that require health and social support. These needs include social and emotional needs (social connections and companionship), civic engagement (meaningfulness and status, the need for having a purpose in later life or being able to contribute usefully to society), healthcare, housing, home modifications and maintenance, domestic assistance, mobility, nutrition and food security, personal care, education (skills development and learning), financial management, respite care, caregiver support, communication (language support or interpreters, information and assistance/referral services) (Abdi et al., 2019; Bedney et al., 2010; Henderson & Caplan, 2008; Jopling, 2020; World Health Organization, 2020).

Support services have been developed to satisfy the needs of older adults and to promote wellbeing and healthy ageing through social networks or relations (Abdi et al., 2019; Jopling, 2020; ten Bruggencate et al., 2019; World Health Organization, 2020).

Social isolation and loneliness may be caused by multiple factors and people respond differently depending on their age and coping skills. It is therefore important to reach out to the older adults to understand their circumstances (the risk factors they are facing and their needs) to be able to provide tailored support for social connections or for accessing services as approaches to reduce social isolation and loneliness (Jopling, 2020; ten Bruggencate et al., 2019).

Digital interventions

Different approaches have been used to reduce social isolation and loneliness including facilitating social connections and providing social support. By providing social support services to meet their needs, opportunities for social connections could be created which could reduce

social isolation and loneliness in older adults. Support for social connections and companionship or for accessing services can be provided through digital technology.

Based on the understanding of the nature and impact of social isolation and loneliness, different strategies have been used in digital interventions to mitigate social isolation and loneliness (Masi et al., 2011). We used these strategies as the typology of intervention categories. Since multiple factors may be involved, multicomponent strategies may also be used to address social isolation and loneliness. The categories include:

- 1) interventions to improve social skills,
- 2) interventions to enhance social interactions,
- 3) interventions to enhance social support,
- 4) interventions for social cognitive training, and
- 5) multicomponent interventions

Mechanisms

The impact of digital interventions can be achieved by four mechanisms of change:

- 1) providing support to building skills for social connections (e.g., Computer and Internet training and use, online university of the third age);
- 2) maintaining existing connections (e.g., video chat with family and friends, personal reminder information and social management system (PRISMS) to engage family and friends in helping receive care);
- 3) creating new connections (e.g., telephone befriending programs, social networking sites, robots and virtual pets, videogames); and
- 4) by changing negative social cognition. (e.g., online cognitive behavioral therapy to teach lonely people to identify and free themselves from negative thoughts and feelings about their relations such as a perception of lack of intimate attachment to their friends or family).

These mechanisms do not map onto the four strategies since some interventions may reduce social isolation or loneliness through more than one mechanism. For example, social networking sites may be used to reduce social isolation and loneliness by maintaining existing connections and by creating new connections. Computer and internet training can be used to maintain connection with family and friends or to create new connections.

Process indicators and outcomes

The effects of interventions depend on how well the interventions were implemented. Process indicators measure activities or outputs that indicate whether the intervention was implemented as planned (Milstein & Wetterhall, 1999). They are preconditions that contribute to the outcomes and are therefore considered proximal indicators of implementation processes or intermediate outcomes (Proctor et al., 2011).

Different levels of outcome measurements for the effects of digital interventions include:

- process indicators – acceptance, adherence, technology use, feasibility, costs, barriers.
- individual level outcomes – loneliness, social isolation, social connectedness, quality of life, anxiety/depression, confidence level, information, communication and technologies (ICT) knowledge and experience, adverse effects.
- community level outcomes – social support, social engagement, social cohesion, social capital, digital divide (disparities in access to technological interventions).

Dimensions

Types of study design

Eligible study designs to be included were completed or on-going systematic reviews, and primary studies with any form of control group, including randomized controlled trials and evaluative quasi-experimental designs with a control group.

We included systematic and scoping reviews based on their PICO question if they explicitly described adequate search methods used to identify studies, eligibility criteria for selection of included studies, methods of critical appraisal of included studies and synthesis or analysis (Moher et al., 2015).

Quasi-experimental design studies were considered eligible if the assignment of participants was based on allocation rules such as alternate assignment (quasi-randomized studies), inclusion of a threshold on a continuous variable (regression discontinuity designs), exogenous variation in the treatment allocation (natural experiments) or other rules including self-selection by investigators or participants, provided data were collected contemporaneously in a comparison group (non-equivalent comparison group design), or an interrupted series design with at least

three data points both before and after a discrete intervention (six-period interrupted time series) (Waddington et al., 2014).

We excluded all studies that used less than six period interrupted time series design, or primary studies without a comparison group design like longitudinal cohort studies with no controls, and cross-sectional studies. We also excluded literature reviews that were not systematic reviews. However, systematic reviews which also include studies without a comparison group design were included.

We did not include qualitative research.

Types of interventions/problem

We considered all types of digital interventions with the aim to reduce social isolation and loneliness. These digital interventions were either one-to-one, or group based. They focused on loneliness, social isolation, or both. We considered any frequency or duration of administration.

We included the following types of digital interventions categorized by strategies.

- [Interventions to improve social skills](#): these are interventions that focus on training in interpersonal social skills such as conversational skills with the aim to enable individuals to form and maintain meaningful relationships. Examples are computer and internet training and use to communicate with others, or online university of the third age which includes courses to facilitate communication with others. We excluded studies that assess computer and internet training for digital literacy and do not assess the use of internet to reduce social isolation or loneliness by focusing on improving social skills.
- [Interventions to enhance social support](#): these are interventions that offer support (e.g., regular contacts, care, or companionship), and guidance in finding and attending new activities or groups. They aim to help individuals make and maintain social connections. Examples include telecare with a component to improve social connections, personal reminder information and social management systems (PRISMS), online support groups and forums, social robots or virtual pets, video games, 3D virtual environments. We excluded studies that assess interventions for care without a communication component or a component to improve connecting with other people e.g., smart home technologies like sensors for monitoring falls, e-health for clinical needs only, online cognitive

behavioral therapy for dementia care only, or online referral systems for healthcare coordination.

- [Interventions to enhance social interactions](#): these are interventions that focus on improving the quality of relationships and increase opportunities for social interactions. They aim to promote connections with family/friends or community and include internet chat facilities, social networking sites, online discussion groups and forums or telephone befriending. Although telephone befrienders could also provide social support, we classified telephone befriending as an intervention to enhance social interactions since the main aim for the service is to connect regularly and build friendship with an older person (Boulton et al., 2020; Gardiner et al., 2018).
- [Interventions for social cognitive training](#): these are interventions that focus on changing negative thinking and feelings about social relationships. They aim to change behaviours, reduce maladaptive cognitions, and increase social connections. Examples include low intensity psychosocial interventions, internet-delivered cognitive behavioral therapy (CBT) or mindfulness interventions.

See Table 1 for categories and other examples

Table 1: Intervention categories

Strategy-based categories and subcategories	Examples
Interventions to improve social skills <ul style="list-style-type: none"> • Skills development • Learning a new activity/language or learning about social skills 	<ul style="list-style-type: none"> • Training in how to use digital technology for communication – e.g., Computer and Internet training and use • Digitally delivered training (e.g., about caregiving/skills building) with a focus on reducing social isolation or loneliness • Digitally delivered learning – e.g., learning a new language, Third age university with courses to facilitate connection with others
Interventions to enhance social interaction <ul style="list-style-type: none"> • Maintain connections • New connections 	<ul style="list-style-type: none"> • Social connections with family/friends – e.g., video chats • Social connections with community – e.g., telephone befriending with volunteers from community

<p>Interventions to enhance social support</p> <ul style="list-style-type: none"> • Healthcare support • Social care support 	<ul style="list-style-type: none"> • Digital/remote ehealth services – e.g., telecare with a component to improve social connections (HomMed Health Telemonitoring system with a communication component) • Digital social and health care coordination with family/friends – e.g., Personal reminder information and social management system (PRISMS) with a communication component • Geolocating/identifying older adults who need services (e.g., Age UK loneliness heat maps) • Socially assistive robots (robopets) and virtual pets • Virtual spaces • Virtual assistants (e.g., Google home, Alexa) • Virtual social support groups • Digital intergenerational approaches • Digital games (e.g., scrabble, chess, cards, exergames) • Digitally delivered activities (e.g., exercise – tai chi, yoga) to mitigate social isolation and loneliness • Digital coordination of health or social care services (e.g., online referrals with a component to improve social connections)
<p>Social cognitive training interventions</p>	<ul style="list-style-type: none"> • Digital cognitive behavioral therapy • Digital mindfulness training • Digital psychoeducation • Digital reminiscence therapy • Digital cognitive behavioral coaching
<p>Multicomponent interventions</p>	<ul style="list-style-type: none"> • Including any of the above in a mixed format (e.g., computer training, messaging, and chat groups)

Comparators were no interventions, other interventions, or usual care.

Type of population (as applicable)

We included older adults, defined as 60 years of age or older (World Health Organization, 2020). If studies included younger and older adults, we included the studies if data could be disaggregated. If data could not be disaggregated, we included studies if the mean age of all participants was at least 65 years of age. To be inclusive, studies or reviews which stated a focus on older adults without providing the age of participants were included.

Types of outcome measures (as applicable)

Outcomes included loneliness and social isolation, as well as other indicators of [social connections](#). Potential harms such as ethical concerns, privacy violations, liability, and cyber-attacks as well as unintended consequences such as increase in social isolation and loneliness, were also included. Community outcomes such as [social support](#), [social engagement](#), [social cohesion](#), [social capital](#), and [digital divide](#) (disparities in access to technological interventions) as well as process indicators (acceptability, adherence, technology use, feasibility, affordability, cost-effectiveness, and barriers), especially for vulnerable populations, were included. (See Table 2.)

Table 2: Outcome categories

Outcomes	Acceptable Measurements
Individual outcomes	
Loneliness	UCLA loneliness scale, de Jong-Gierveld loneliness scale, other scales e.g., Social and Emotional Loneliness Scale, Hughes loneliness scale
Social isolation	Lubben’s Social Network Scale, Social Network Index, PROMIS social isolation 6-I scale
Social connectedness/interactions/networks or life satisfaction	Lee and Robin’s Social Connectedness Scale; Number of contacts; Frequency of social interactions; Satisfaction with interaction; Index of support satisfaction; Support network satisfaction; Companionship scale satisfaction
Social support	Duke-UNC Functional Social Support Questionnaire; Social support scale by Schuster et al; Hsiung’s Social Support Behaviors Scale; Family and Friendship Contacts Scale; Personal Resource Questionnaire; Interpersonal Support Evaluation List (ISEL); e-Diabetes Social Support Scale; a bespoke six-item scale measuring women’s perception of emotional and instrumental support
Well-being/Quality of life	MOS SF-36 Health Survey; Work and Social Adjustment Scale (WSAS); WHOQOL
Anxiety/depression	Beck Depression Inventory (BDI); Depression Adjective Check List (DACL) Form E; Geriatric depression scale; The Centre for Epidemiological Studies Depression Scale (CES-D)
Confidence level	Rosenberg Self-Esteem Scale

Information, communication and technologies (ICT) knowledge and experience	Questionnaire
Adverse effects	Privacy violations, liability, cyber-attacks, negative effect on well-being from emotional attachment to devices, increases in social isolation and loneliness
Community outcomes	
Social support	Duke-UNC Functional Social Support Questionnaire, Social support scale, social Provisions scale
Social engagement	Engagement in Meaningful Activities Survey (EMAS)
Social cohesion (measures ability to trust and expect reciprocity from neighbors)	The Group Cohesion Scale-Revised; Group Therapy Experience Scale, Group Environment Questionnaire
Social capital (measures trustworthy and reciprocal social networks that people can access in their community)	The World Bank's integrated questionnaire for the measurement of social capital (SC-IQ)
Digital divide	Lack of affordability/access to technology, lack of affordability/access to broadband or Wi-Fi, data poverty, lack of digital skills or confidence to access services and support online
Process indicators	
Acceptability (technology adoption)	Various survey tools to measure acceptability
Adherence (training adherence)	Various survey tools to measure adherence
Technology use	Frequency of use
Feasibility	Various survey tools to measure feasibility
Affordability	Various survey tools to measure affordability
Cost-effectiveness	Cost-effectiveness analysis
Barriers	e.g., language and cultural barriers, financial accessibility, hearing or vision impairments, personal barriers such as dislike of robotpets, digital literacy, lack of familiarity with digital technologies, Lack of confidence in using digital technologies

Outcomes were not used as eligibility criteria. However, eligible studies and systematic reviews did have a focus on social isolation and loneliness. Studies and reviews assessing interventions with a stated aim to reduce social isolation and loneliness were eligible. Those that assessed the effects of interventions on social isolation and/or loneliness as a primary outcome or considered

other indicators of social connections including quality of life, anxiety/depression, social support, social engagement, social cohesion, and social capital were also included.

Studies and reviews assessing the effect of interventions on anxiety or depression with a focus on mental health rather than social isolation or loneliness were excluded.

Other eligibility criteria

Types of Location/Situation (as applicable)

We included all country settings as defined by the World Health Organization regions (African Region, Region of the Americas, South-East Asian Region, European Region, Eastern Mediterranean Region, Western Pacific Region) (World Health Organization, 2019) and the World Bank classification by income: low income economies, lower-middle income economies, upper-middle income economies, high income economies (World Bank 2021).

Primary studies and systematic reviews that did not report the countries were not excluded.

Types of settings (as applicable)

All settings except hospital settings were included, that is, people living in supportive care institutions (nursing home or long-term care and assisted living facilities) and in the community (residential or personal home).

Search methods and sources

We designed a search strategy with an information scientist (DS) in consultation with Tomas Allen (WHO information specialist). We searched the following databases from inception to May 16, 2021, with no language restrictions. Ovid MEDLINE, Embase, APA PsycInfo via Ovid, CINAHL via EBSCO, Web of Science via Clarivate, ProQuest (all databases), International Bibliography of the Social Sciences (IBSS) via ProQuest, EBSCO (all databases except CINAHL), Global Index Medicus, and Epistemonikos. The full search strategies are in [Appendix 2](#).

We screened reference lists of all included systematic reviews in Eppi-Reviewer to identify additional studies. We also contacted stakeholders for information about ongoing studies.

Analysis and presentation


Report structure

The report has the standard sections: abstract, plain language summary, background, methods, results, discussion, and conclusion.

The report includes the flow of studies, included studies, excluded studies and any studies awaiting assessment, as well as synthesis of included studies. We presented the PRISMA flowchart and conceptual framework. We also included tables and figures that provide a summary of the distribution of primary studies and systematic reviews across the coding categories such as the type of studies, quality of the systematic reviews, types of interventions, needs, types of populations, outcomes, settings, and geographic distribution.

The evidence and gap map has interventions as the row dimension and outcomes as the column dimension. Bubbles of different sizes represent included studies and different colors are used to identify the primary studies and methodological quality of the systematic reviews. The filters used in the map depend on the number of included studies and coded information. See a sample of the map in Figure 4.

Figure 4: Sample map



Filters for presentation

Additional dimensions of interest that were used as filters included the publication status of included studies, study design, World Bank classification by income (low income economies, lower-middle income economies, upper-middle income economies, high income economies), and WHO regions (African Region, Region of the Americas, South-East Asian Region, European Region, Eastern Mediterranean Region, Western Pacific Region), setting (personal home, independent living/residential home, assisted living, long-term care/nursing home), health status/condition.

We documented which needs beyond mitigating social isolation and/or loneliness of older adults are being met by digital interventions, using a framework developed from our citizen and stakeholder engagement consultation, which includes social and emotional needs, civic engagement and social participation, healthcare, housing, home modifications and maintenance, domestic assistance, mobility, nutrition and food security, personal care, education, financial management, respite care, caregiver support, communication.

We documented the focus of the intervention as aimed at social isolation, loneliness, or both. We also had filters for the intervention format, how technologies are used, as well as how training on how to use digital technology was delivered.

Equity analysis

We documented whether studies were focused on populations who are at risk or experiencing barriers to health and social care or health inequities across age, sex, ethnicity, income, or other factors. We used the PROGRESS-Plus acronym to describe factors associated with health inequities. For these studies, we documented how potentially vulnerable older adults are defined and identified (e.g., using case finding, outreach, screening).

In addition, for each study, we assessed whether studies have analyzed differences in effects for populations experiencing inequities, using the PROGRESS factors (Place of residence (urban/rural), Race/ethnicity/culture and language, Occupation, Gender or sex, Religion, Education, Socioeconomic status, Social capital. We also assessed analysis across additional “Plus” factors which are known to be important for older adults, including age, disability, social frailty, health status, being socially isolated or at risk, being lonely or at risk, living situation, and digital literacy.

Dependency

Multiple reports of the same study were treated as one study. A study with multiple interventions or outcomes was shown multiple times on the map (for each intervention or outcome identified). Systematic reviews were mapped to the interventions and outcomes as defined by the question of the systematic review. Primary studies that met the eligibility criteria were mapped as well regardless of whether they are included in one or more systematic reviews.

Data collection and analysis

Screening and study selection

Titles and abstracts and full text of potentially eligible articles were screened independently following the eligibility criteria in duplicate (by VW, EG, VB, PG, TH, SA, NE, JE, HW and OD) using the Eppi-Reviewer web-based software program (Thomas et al., 2020). We screened systematic reviews based on their PICO questions. Disagreements were resolved by discussion. See [Appendix 3](#) for detailed eligibility criteria.

We used machine learning text mining to support screening at the title and abstract stage. After screening approximately 10 per cent of the titles and abstracts, we used the priority screening function which developed a classifier based on the probability of inclusion determined from the preliminary screening results. We however, double screened all the search results to ensure all potentially eligible studies were captured for the full text screening stage.

We also screened reference lists of eligible systematic reviews to identify additional studies.

Data extraction and management

We developed and pilot tested a data extraction code set in Eppi-Reviewer for data collection (see extraction code in [Appendix 4](#)). After the pilot test, members of the team (EG, VB, TH, EB, AW, AA, and SD) individually extracted and coded data. Automation and text mining were not used for coding.

The coding categories included study characteristics (study design, publication status, methodological quality assessment of systematic reviews), intervention categories and subcategories, intervention focus (loneliness, social isolation, or both), intervention format, how technologies are used, how training on how to use digital technology was delivered,

outcome domains and subdomains, population characteristics, needs, setting, and location (countries, World Health Organization region, and World Bank classification by income).

We coded the location (country) if it was reported for the primary study or the included studies for reviews. If a review had multiple studies conducted in the same country, it received only one code for the country. For example, a review with six included studies conducted in the USA and two studies conducted in Australia, had only two codes – one for USA and one for Australia.

We coded description of the population characteristics using the PROGRESS-Plus framework, defined as Place of residence (urban/rural), Race/ethnicity/culture and language, Occupation, Gender/sex, Religion, Education, Socioeconomic status, Social capital(marital status) and additional (plus) factors such as age groups, health status/condition, frailty, disability, living situations, digital literacy, social isolation, and loneliness.

We considered how the study population was selected based on whether they are disadvantaged across any PROGRESS-Plus factors.

We also coded whether there was any analysis that aimed to understand potential differences across any PROGRESS-Plus factors.

Given the expected size of the map (over 200 studies), we did not contact organizations or authors of studies and systematic reviews for missing information.

Tools for assessing risk of bias/study quality of included reviews

We assessed the methodological quality of systematic reviews in duplicate (by EG, SD, VB, TH, NE and AW) using the AMSTAR 2 tool (Shea et al., 2017). Any disagreements were resolved by discussion. As per guidance for evidence maps, primary studies were not assessed for risk of bias or methodological quality (Snilstveit et al., 2016; White et al., 2020). A modified AMSTAR2 assessment was conducted for 20 scoping reviews; we did not assess items about risk of bias of included studies since it is optional for scoping reviews (Peters et al., 2021).

Methods for mapping

We used the EPPI-Mapping tool (Digital Solution Foundry and EPPI_Centre, 2020) to develop the evidence and gap map.


Results

Description of studies

Results of the search

Our search retrieved 15,324 records from databases and 1829 records from included systematic reviews. After duplicate records were removed, 11974 articles were screened by title and abstract in duplicate. From this, 633 articles were assessed for eligibility, in duplicate. Two hundred articles were included for the purpose of this EGM. See Figure 5 for the PRISMA flow diagram.

Figure 5: PRISMA flow diagram



Excluded studies

A total of 433 studies were excluded at the full-text review phase due to inappropriate evidence/aim ($n = 175$), inappropriate intervention ($n = 58$), inappropriate study design ($n = 96$), or inappropriate target population ($n = 97$). See figure 5 for the PRISMA flow diagram.

See key excluded studies in the table for [Characteristics of excluded studies](#). Three studies (Bolle et al., 2015; Clarkson et al., 2018; Sumner et al., 2021) were excluded because the aim was not to reduce social isolation or loneliness. Bolle et al., 2015 assessed online health information tools to improve health outcomes. Clarkson et al., 2018 assessed home support interventions to

inform dementia care and Sumner et al., 2021 assessed co-designed interventions to support ageing in place.

Six studies did not assess digital interventions (Dickens et al., 2011; Giné-Garriga et al., 2017; Jones et al., 2019; Nicholson, 2012; Toh et al., 2016; Zeppego et al., 2019).

Eight studies were excluded for wrong study design; five (Fan, 2016; Forsman & Nordmyr, 2017; Gorenko et al., 2021; Selak et al., 2019; Winterton & Warburton, 2011) were not systematic reviews of effectiveness and three were primary studies with no control groups (Perkins & LaMartin, 2012; Preston & Moore, 2019; Rebollar & Francisco, 2015).

Six studies (Burkow et al., 2015; Cooper et al., 2014; Dam et al., 2017; Erfani & Abedin, 2018; Lara et al., 2016; Nijman et al., 2019) included people less than 60 years old and did not provide disaggregated data for people 60 years and older.

Synthesis of included studies

We included 200 articles that utilized digital interventions to reduce social isolation and/or loneliness; 103 were primary studies and 97 were systematic reviews. See Supporting Information interactive EGM <https://globalageing.cochrane.org/cochrane-reviews>.

The primary studies included both randomized controlled trials (RCTs) (n = 81) and non-randomized studies (NRSIs) (n = 22). The systematic reviews included both systematic reviews (n = 77) and scoping reviews (n= 20) that explored the effectiveness of interventions to reduce social isolation and loneliness. Most of the included publications were completed (n = 170, 78 primary studies and 92 reviews) while others were ongoing registered trials and protocols (n = 25, 21 primary studies and 4 reviews). There were five conference abstracts included (n = 5, 4 primary studies and 1 review). See Table 3 (Characteristics of included articles).

Table 3: Characteristics of included articles

Characteristics of included articles		Number of articles
Study design	Primary studies	103
	- randomized studies	81
	- non-randomized studies	22


	Reviews	97
	- systematic reviews	77
	- scoping reviews	20
Publication status	Completed	78 primary studies / 92 reviews
	On-going	21 primary studies / 4 reviews
	Conference abstracts	4 primary studies / 1 review
Intervention focus	Social isolation	34 primary studies / 54 reviews
	Loneliness	41 primary studies / 23 reviews
	Both social isolation and loneliness	34 primary studies / 54 reviews
Intervention format	One-on-one	29 primary studies / 10 reviews
	Group-based	56 primary studies / 23 reviews
	One-on-one and group based	11 primary studies / 28 reviews
	Unspecified	7 primary studies / 26 reviews
Intervention strategies to reduce social isolation and loneliness	Social cognitive training	21 primary studies / 19 reviews
	Improving social skills	37 primary studies / 40 reviews
	Enhancing social interactions	30 primary studies / 58 reviews
	Enhancing social support	49 primary studies / 81 reviews
	Multicomponent	16 primary studies / 3 reviews
Outcomes	Individual level outcomes	95 primary studies / 96 reviews
	Community level outcomes	40 primary studies / 40 reviews
	Process indicators	28 primary studies / 31 reviews
Most reported population sociodemographic characteristics	Gender/sex	83 primary studies / 78 reviews
	Age range (70-80 years old)	78 primary studies / 55 reviews
	Education	50 primary studies / 6 reviews
	Health condition (dementia)	16 primary studies / 30 reviews

Settings	Personal home	36 primary studies / 45 reviews
	Long-term care (nursing homes)	22 primary studies / 45 reviews
	Independent living (residential homes)	12 primary studies / 23 reviews
	Assisted living	5 primary studies / 10 reviews
Most common locations for included studies	USA	42 primary studies / 44 reviews
	The Netherlands	8 primary studies / 25 reviews
	UK	6 primary studies / 21 reviews
	Canada	6 primary studies / 16 reviews
	Australia	5 primary studies / 22 reviews

Interventions


The intervention focus for included publications was on addressing social isolation and loneliness together (n = 88, 34 primary studies and 54 reviews), or on loneliness (n = 64, 41 primary studies and 23 reviews) or social isolation (n = 48, 28 primary studies and 20 reviews) alone (Figure 6).

Figure 6: Intervention focus




Included publications reported delivery of interventions as one-on-one (n = 79, 56 primary studies and 23 reviews), group-based (n = 39, 29 primary studies and 10 reviews), or both one-on-one and group-based (n = 49, 11 primary studies and 38 reviews). The mode of delivery was unspecified in some publications (n = 33, 7 primary studies and 26 reviews) (Figure 7).

Figure 7: Intervention format




Interventions were categorized based on their strategies to reduce social isolation or loneliness: interventions for social cognitive training (n = 40, 21 primary studies and 19 reviews), improving social skills (n = 77, 37 primary studies and 40 reviews), enhancing social interactions (n = 88, 30 primary studies and 58 reviews), enhancing social support (n = 130, 49 primary studies and 81 reviews), and multicomponent interventions (n = 19, 16 primary studies and 3 reviews) (Figure 8).

Figure 8: Intervention strategies to reduce social isolation and loneliness



Each coded intervention category included multiple subcategories. Interventions for social cognitive training consisted of five subcategories: digital cognitive behavioral therapy (n = 15, 7 primary studies and 8 reviews), digital psychoeducation (n = 12, 5 primary studies and 7 reviews), digital cognitive behavioral coaching (n = 10, 8 primary studies and 2 reviews), digital reminiscence therapy (n = 6, 2 primary studies and 4 reviews), and digital mindfulness training (n = 4, 3 primary studies and 1 review) (Figure 9).


Figure 9: Interventions for social cognitive training



Cognitive behavioral therapy approaches help individuals recognize and change negative thinking whereas cognitive behavioral coaching approaches help individuals reach their desired goals. Reminiscence therapy involves recalling past events and encourages older adults to communicate and interact with someone in the present. Mindfulness training teaches people to consciously pay attention to their thoughts and feelings and to disengage from negative thoughts, and unhealthy habits that may render them vulnerable. Psychoeducation therapy develops knowledge and understanding of psychological conditions to help individuals cope with them.


Interventions for improving social skills had three subcategories: training in how to use digital technology for communication was most frequent (n = 55, 28 primary studies and 27 reviews), followed by digitally delivered training (n = 23, 9 primary studies and 14 reviews), and no studies or reviews for digitally delivered learning (Figure 10). Training in how to use digital technology was mainly about computer and internet training for communication with family and friends. Half of the primary studies and reviews on digitally delivered training interventions were technology-based interventions for caregivers of older adults with dementia.

Figure 10: Interventions for improving social skills



The interventions for enhancing social interactions comprised two subcategories: digital connections with family/friends (n = 57, 13 primary studies and 44 reviews), and digital connections with community including healthcare and social workers and volunteers (n = 61, 22 primary studies and 39 reviews) (Figure 11).

Figure 11: Interventions for enhancing social interactions



Interventions for enhancing social support had 11 subcategories with most studies involving the following six subcategories: socially assistive robots and virtual pets (n = 51, 10 primary studies and 41 reviews), digital games (n = 29, 9 primary studies and 20 reviews), virtual spaces (n = 28, 11 primary studies and 16 reviews), digital/remote e-health services that also provided companionship or social connections (n = 28, 6 primary studies and 22 reviews), virtual social support groups (n = 27, 10 primary studies and 17 reviews) and digitally delivered activities, mostly physical exercise and exergaming activities (n = 19, 9 primary studies and 20 reviews). Few studies and reviews assessed digital social and healthcare coordination with family/friends (n = 5, 1 primary study and 4 reviews), geolocating/identifying older adults who need services (n = 2 reviews), virtual assistants (n = 3, 1 primary study and 2 reviews), digital intergenerational approaches (n = 3 reviews), and digital coordination of health or social care services (n = 4, 2 primary studies and 2 reviews) (Table 4).

Table 4: Interventions for enhancing social support


Interventions for enhancing social support	Primary studies (n = 103)	Reviews (n = 97)
Geolocating/Identifying older adults who need services	0	2
Digital intergenerational approaches	0	3
Virtual assistants	1	2
Digital social and health care coordination with family/friends	1	4
Digital coordination of health or social care services	2	2
Digital/Remote e-health services	6	22
Digitally delivered activities	9	10
Digital games	9	20
Virtual social support groups	10	17

Socially assistive robots and virtual pets	10	41
Virtual spaces	12	16

Outcomes


Three outcome categories were used to code publications: individual level outcomes (n = 191, 95 primary studies and 96 reviews), community level outcomes (n = 80, 40 primary studies and 40 reviews), and process indicators (n = 59, 28 primary studies and 31 reviews) (Figure 12).

Figure 12: Outcome categories



Many publications reported multiple outcomes and all outcomes of interest were coded. Individual level outcomes included: loneliness (n = 123, 56 primary studies and 67 reviews), anxiety/depression (n = 112, 57 primary studies and 55 reviews), quality of life/wellbeing (n = 90, 41 primary studies and 49 reviews), social connectedness (n = 54, 17 primary studies and 37 reviews), social isolation (n = 52, 15 primary studies and 37 reviews), confidence level or self-esteem (n = 23, 8 primary studies and 15 reviews), and adverse effects (n = 8, 2 primary studies and 6 reviews) (Figure 13).

Figure 13: Individual level outcomes




Adverse effects were mostly privacy and ethical issues reported in three reviews, excessive attachment to robopets with detrimental effects in one review, aggravated musculoskeletal symptoms and fall risks from exergaming in one review, negative impact of adverse media in social networking sites reported in one review, and increase in loneliness in one review. The two primary studies were on-going and planned to measure adverse effects but didn't indicate which ones were anticipated.

Community level outcomes consisted of the following: social support defined as the actual or perceived availability of resources (e.g., tangible, informational, emotional help) from others, typically one's social network (n = 60, 31 primary studies and 29 reviews), social engagement which reflects participation in meaningful activities with others (n = 21, 8 primary studies and 13 reviews), social capital which refers to an appraisal of the social resources or networks people can access in their community (n = 5, 1 primary study and 4 reviews), social cohesion, a measure of mutual community trust and solidarity (n = 1 review). Digital divide, defined as disparities in access to technological interventions (e.g., smartphones, computers, and the internet) which may be due to lack of affordability or access to technology, broadband or Wi-Fi, data poverty or


geographic location, lack of digital skills and the confidence to access online services and support (n = 0) was not measured in any of the included articles (Figure 14). However, one review discussed the barriers caused by digital divide and three primary studies and three reviews each discussed the value of accessibility of digital interventions.

Figure 14: Community level outcomes



Lastly, process indicators consisted of seven outcomes: acceptance (n = 26, 11 primary studies and 14 reviews), technology use (n = 19, 6 primary studies and 13 reviews), feasibility (n = 15, 6 primary studies and 9 reviews), adherence (n = 14, 10 primary studies and 4 reviews), cost-effectiveness (n = 9, 1 primary study and 8 reviews), barriers (n = 4 reviews), and affordability (n = 0) which was not measured in any of the included studies or reviews. (Figure 15).


Figure 15: Process indicators



Risk of bias in included reviews

Most reviews were classified as critically low quality on the AMSTAR2 Quality Assessment tool and results were as follows: critically low (n = 70), low (n = 15), moderate (n = 6), and high (n = 2) quality. Protocols (n = 4) were not assessed (Figure 16).

Figure 16: Quality assessment of reviews



Additional dimensions

How Technologies Are Used

Technology was used in various ways, and multiple methods were considered in some publications. The most utilized method was videoconferencing (n = 65, 23 primary studies and 42 reviews), followed by telephone calls (n = 55, 20 primary studies and 35 reviews) and robots or virtual companions (n = 51, 10 primary studies and 41 reviews). Mobility tools (n = 1 review), listening to or creating music (n = 3 reviews) and virtual assistants (n = 4, 1 primary study and 3 reviews) were among the least ways in which technology was utilized, and 10 publications (5 primary studies and 5 reviews) did not specify how technologies were used (Table 5).

Table 5: How technologies are used

How technologies are used	Primary studies (n = 103)	Reviews (n = 97)
Mobility tools	0	1
Listening to or creating Music	0	3

Virtual assistants	1	3
Discussion forums	3	11
Participating in an activity	15	6
Social networking sites	12	16
Digital games	10	20
Messaging	14	17
E-mail	13	22
Virtual spaces or classrooms with messaging capabilities	16	28
Robots and virtual companions	10	41
Telephone calls*	20	35
Videoconferencing	23	42
Unspecified	5	5


* Smartphones were specified in 32 of 55 primary studies and systematic reviews. For this EGM, we considered all telephone calls as a strategy for remotely delivered interventions.

Region

Publications included in this EGM come from different countries worldwide. We coded the countries where primary studies were conducted. For reviews, if multiple included studies were conducted in the same country, we coded the country just once. However, if primary studies were included in multiple reviews, the countries where they were conducted were coded multiple times (i.e., once for each review).


Many of the included publications had at least one study conducted in the USA (n = 86, 42 primary studies and 44 reviews) followed by the Netherlands (n = 33, 8 primary studies and 25 reviews), UK (n = 27, 6 primary studies and 21 reviews), Canada (n = 22, 6 primary studies and 16 reviews) and Australia (n = 27, 5 primary studies and 22 reviews). (See geographic heat map for primary studies in Figure 17).

Figure 17: Geographic heat map for primary studies




In terms of classification by WHO regions (Figure 18) most publications were from the Americas (n = 95, 48 primary studies and 47 reviews), the European Region (n = 77, 33 primary studies and 44 reviews) or Western Pacific Region (n = 58, 20 primary studies and 38 reviews), while very few were from the South-East Asian (n = 5, 1 primary study and 4 reviews), African (n = 3, 1 primary study and 2 reviews), and Eastern Mediterranean Regions (n = 2 primary studies). Forty reviews did not specify the country where included primary studies were conducted, thus the WHO region and World Bank income region, could not be determined.

Figure 18: World Health Organization regions



Following the World Bank income classification, most publications were from high-income (n = 146, 92 primary studies and 54 reviews) or upper-middle-income countries (n = 22, 10 primary and 12 reviews), with none from low-income countries (Figure 19).


Figure 19: World Bank income classification



Settings

The setting for most interventions described was the personal home of participants (n = 81, 36 primary studies and 45 reviews), followed by long-term care/nursing homes (n = 67, 22 primary studies and 45 reviews). Few took place in independent living (n = 35, 12 primary studies and 23 reviews) or assisted living settings (n = 15, 5 primary studies and 10 reviews). Again, some publications included more than one setting for the intervention, and all reported settings were coded as such. Sixty-eight publications (35 primary studies and 33 reviews) did not specify the setting (Figure 20).

Figure 20: Setting



Participant needs

Different types of participant needs beyond social isolation and loneliness were identified. Social and emotional needs (n = 191, 97 primary studies and 94 reviews) were the most identified, followed by clinical/health needs (n = 19, 5 primary studies and 14 reviews) and caregiver support (n = 22, 8 primary studies and 14 reviews). Some publications reported interventions targeting more than one need, while no included article targeted participants' accommodation or financial assistance needs (Table 6).

Table 6: Participant needs


Participant needs	Primary studies (n = 103)	Reviews (n = 97)

Social and emotional needs	97	94
Caregiver support	8	14
Clinical/Health needs	5	14
Mobility	7	4
Purpose in Life	5	2
Personal care needs	1	5
Skills development	2	3
Meals	2	0
Respite care	1	2
Communication (language support/interpreters)	1	1
Domestic assistance	1	1
Care navigation support or task orientation	1	0
Learning	0	5
Accommodation	0	0
Financial assistance	0	0

Participant Characteristics (PROGRESS-Plus)

Every PROGRESS-Plus factor was used to describe participants in at least one included article (Figure 21). The PROGRESS acronym stands for Place of residence, Race/ethnicity, Occupation, Gender or sex, Religion, Education, Socioeconomic status, Social capital (marital status). For the purpose of this EGM, we also considered living alone, age group, disability, health condition, and being caregivers as plus factors.

Figure 21: PROGRESS-Plus categories used to describe participants




Gender or sex (n = 161, 83 primary studies and 78 reviews) and age (n = 140, 84 primary studies and 56 reviews) were the most reported characteristics, while occupation (n = 8, 7 primary studies and 1 review), and religion (n=1 primary study) were the least reported.

Of the 161 articles specifying gender or sex, most included male and female participants. Two reviews included some studies with only male participants and 8 reviews included some studies with only female participants; 4 primary studies included only female participants, and gender or sex was unspecified in 39 articles (20 primary studies and 19 reviews). No articles included individuals from the LGBTQIA2S+ (lesbian, gay, bisexual, transgender, queer (or sometimes questioning), and two-spirited) community.

Of the articles that reported age, the 70–80-year-old range was most reported (n =133, 78 primary studies and 55 reviews), and the age range restricted to older than 80 was least reported (n = 1 primary study). Age of participants was not specified in 67 articles (19 primary studies and 48 reviews) (Figure 22).


Figure 22: Participant age groups



Health condition of participants

The health condition of participants was reported in 75 articles (33 primary studies and 42 reviews). Dementia was most common (n = 46, 16 primary studies and 30 reviews), followed by other non-communicable diseases (n = 13, 8 primary studies and 5 reviews). Included articles did not report on communicable diseases, discharge from hospital, and end-of-life/palliative care (Figure 23).


Figure 23: Health conditions of participants



Equity Analysis

Of the 200 included articles, 78 primary studies and 52 reviews focused on recruiting older adults vulnerable across one or more PROGRESS-Plus factors. (Figure 24). Most included articles focused on living situation i.e., older adults living alone or in a nursing home (n = 50, 30 primary studies and 20 reviews), health status (including dementia, depression and other chronic conditions) of older adults (n = 47, 24 primary studies and 23 reviews), socially isolated older adults or at risk of social isolation (n = 20, 13 primary studies and 7 reviews), or lonely older adults or at risk of loneliness (n = 19, 11 primary studies and 8 reviews). Few articles focused on other equity relevant factors including occupation (n = 9, 6 primary studies and 3 reviews), frailty (n = 7, 5 primary studies and 2 reviews), ethnicity (n = 4 primary studies), place of residence, urban or rural (n = 4 primary studies), gender/sex (n = 3 primary studies), digital literacy (n = 3, 2 primary studies and 1 review), disability (n = 2 primary studies), socioeconomic status (n = 2 primary studies), education level (n = 1 primary study), or social capital (n = 1 primary study). No articles focused on religion.


Figure 24: Equity focus of vulnerability across PROGRESS-Plus factors



Very few studies focused on multiple vulnerability categories, for example, frail and socially isolated older adults (n = 1), rural isolated family caregivers (n = 1), Chinese socially isolated immigrants (n = 1), low-income women living alone (n = 1), populations at risk of both social isolation and loneliness (n = 9).


Very few studies described how at-risk populations were recruited (Figure 25). They were identified through outreach (n = 11, 10 primary studies and 1 review), community-based programs (n = 11, 8 primary studies and 3 reviews), case finding (n = 7, 6 primary studies and 1 review), or through screening in primary care (n = 3 primary studies). One primary study used all four approaches to identify at-risk populations.

Figure 25: Recruitment strategies for at-risk populations



We also assessed whether studies analyzed differences in effects for populations experiencing inequities across PROGRESS-Plus categories. Despite the large number of articles that considered at least one equity focus, only five articles (4 primary studies and 1 review) assessed the differences in effects across any PROGRESS-Plus factor including gender or sex (n = 1 study), education level (n = 3 primary studies), age (n = 3 primary studies), living situation (n = 1 primary study), and health status (n = 1 review). No equity analyses were completed for place of residence, race/ethnicity, occupation, religion, socioeconomic status, social capital, frailty, digital literacy, being socially isolated or at risk, being lonely or at risk, or disability (Figure 26).

Figure 26: Equity analyses to assess differences in effects across any PROGRESS-Plus factor



Stakeholder feedback on framework and draft map

We surveyed 30 people and got 10 (33%) responses. Eight of the 10 (80%) people found that the intervention categories made sense, one (1%) person was neutral and one (1%) disagreed with the category social skills training. We decided to keep social skills training based on the Masi framework (Masi et al., 2011).

Six (60%) people found the outcome categories made sense, three (30%) were neutral and one (1%) disagreed with including depression/anxiety, quality of life and confidence level as outcomes. We kept these outcomes because they were indicators of social connections (Elder & Retrum, 2012; Windle et al., 2012) and were reported in many included articles – 112/200 studies measured anxiety/depression, 90/200 studies measured quality of life, and 23/200 studies measured confidence. Although six (60%) people easily found the information they were looking for on the draft map and one (16.67%) person was neutral, they provided helpful suggestions for knowledge translation activities and improving the usability of the map such as creating a short video of how to use the map.

Discussion

Summary of main results

This evidence and gap map has 200 included articles and there are almost as many primary studies (n = 103) as reviews (n = 97), with most conducted in the Americas, the European and Western Pacific regions. The high ratio of reviews to primary studies indicates a surge in research with over 25% of the reviews conducted since the COVID-19 pandemic but most reviews are of critically low quality.

The evidence is unevenly distributed across intervention and outcome subcategories with a similar trend across primary studies and reviews. The majority of included primary studies and reviews assessed digital interventions to enhance social interactions with family and friends and the community via videoconferencing and telephone calls. Digital interventions to enhance social support, particularly socially assistive robots, and virtual pets were the next most assessed subcategory of interventions. The social cognitive training strategy was the least assessed of the four intervention strategies although it is considered the most effective (Mann et al., 2017; Masi et al., 2011).

The most assessed outcomes were reducing loneliness and depression and improving the quality of life of older adults. Only six reviews reported adverse events and two on-going primary studies planned to report adverse effects. The most reported age range was 70-80 years old and most common needs addressed were social and emotional needs.

Although many primary studies and reviews focused on recruiting older adults vulnerable across one or more PROGRESS-Plus factors, very few described how at-risk populations were recruited or conducted any equity analysis to assess differences in effects for populations experiencing inequities across PROGRESS-Plus categories.

Areas of major gaps in the evidence

Major gaps were identified in outcome measures including adverse effects, community level outcomes and process indicators. Only six reviews reported adverse effects. No included studies or reviews assessed affordability or digital divide, which refers to disparities in access to technological interventions, although three primary studies and three reviews discussed the value of accessibility of digital interventions and one of the reviews discussed the barriers caused by digital divide. There were no primary studies or reviews conducted in low-income

countries. Only one primary study restricted participants to 80 years and older and no study or review included participants from the LGBTQIA2S+ community.

Potential biases in the mapping process

The search strategy was comprehensive and covered published and grey literature until 16 May 2021. We used rigorous methods including duplicate screening of all the identified records and reference lists of included reviews and assessment of methodological quality of reviews. Since our research question is about the effectiveness of digital interventions for social isolation or loneliness, we included only study designs that are appropriate for assessing effectiveness. We did not use outcomes as eligibility criteria. We included studies that assessed the effects of digital interventions on social isolation and loneliness as well as other indicators of social connections such as quality of life, depression.

With the growing research in this area, we may have missed more recent primary studies and reviews. Although there were no language restrictions applied, we may have missed studies and reviews published in non-English language since mainly English databases were searched.

Mapping the interventions into mutually exclusive categories in the evidence and gap map was challenging due to the lack of consistent terminology and a standardized framework for the classification of interventions for social isolation and loneliness.

Limitations of the EGM

The map does not include evidence from studies with no control groups or qualitative research. Although these studies might provide useful information about digital interventions for social isolation and loneliness in older adults, they do not assess the effectiveness of interventions.

Most studies and reviews do not focus on reaching the most vulnerable older adults including specific groups such as lonely/socially isolated or at-risk, low socio-economic status, low literacy, ethnic minority groups, LGBTQIA2S+ community or populations in low-income countries where access to digital technology is limited. No included studies or reviews assessed affordability or accessibility of digital interventions, but three primary studies and three reviews discussed the value of accessibility of digital interventions and one of the reviews discussed the barriers caused by digital divide.

Stakeholder Engagement throughout the EGM process

Citizens were involved in the development of this EGM, and they provided constructive and valuable feedback that was incorporated in the process. Only 29% of stakeholders, including citizens, provided feedback on the framework and draft map.

Although one stakeholder disagreed with social skills training as an intervention category, we decided to keep it since it is included in the Masi framework as interventions that focused on improving interpersonal communication skills (Masi et al., 2011). One stakeholder disagreed with including depression/anxiety, quality of life and confidence level as outcomes but these outcomes have been described as indicators of social connections (Elder & Retrum, 2012; Windle et al., 2012) and were reported in many included articles.

Authors' conclusions

Implications for research, practice and/or policy

Social isolation and loneliness are growing public health concerns, which have been highlighted during the Covid-19 pandemic. This evidence and gap map shows available evidence on effectiveness of digital interventions for reducing social isolation or loneliness in older adults. Over 70% of the systematic reviews have critically low quality, 2% have high quality, and 25% have been published since the pandemic. Clusters of evidence of critically low quality exist mainly for digital interventions to enhance social interactions with family and friends and the community via videoconferencing and telephone calls as well as digital interventions to enhance social support, particularly socially assistive robots, and virtual pets. Most of the digital interventions were focused on reducing loneliness and depression and improving quality of life of older adults.

Major gaps were lack of evidence on equity analysis, community level outcomes and process indicators. Affordability and the digital divide were not assessed in any of the included studies or reviews although barriers caused by the digital divide may increase health inequities and have been key challenges for organizations and governments concerning digital interventions during the COVID-19 pandemic restrictions. Many studies did not focus on reaching the most vulnerable population groups. The evidence is predominantly from high-income countries with none in low-income countries. Other challenges faced by organizations and governments are related to the adverse effects of digital interventions such as privacy and ethical issues. However, very few primary studies and reviews have assessed adverse effects.

This evidence and gap map is the starting point for building the evidence architecture and will guide a future research agenda to standardize definitions of concepts, intervention

characteristics, and outcome measurements as well as identify which digital interventions are effective for reducing social isolation and loneliness in older adults taking into consideration their specific needs and contexts.

Although there is proliferation of evidence in recent years, it is unevenly distributed, and the systematic reviews are of poor quality highlighting the need for high quality research.

Registration of reviews may help to reduce duplication. This map can guide researchers and funders to consider areas of major gaps as priorities for further research.

Acknowledgements

We acknowledge the contribution of Salman Abdisalam, , Priya Grewal, , Jinane ElHage, , Harry Wang, and Omar Dewidar in screening and coding of articles as well as Tomas Allen for reviewing the search strategies.

We gratefully acknowledge the participation of Rheda Adekpedjou and stakeholders in our stakeholder consultation and citizen focus groups.

This project was funded by the World Health Organization.

Contributions of authors

- Content: PH, CM, VW, EG, RS, SB, NH, EB, DK
- EGM methods: VW, EG, VB, EB, DK, SD, RD, NE, TH, AW, EBev, PD, JH
- Information retrieval: DS

All authors read and approved the review.

Declarations of interest

Vivian Welch is editor in chief of the Campbell Collaboration. The editorial process was handled by an independent editor and VW had no input in the editorial process or decisions.

Elisabeth Boulton and Dylan Kneale are joint lead authors of a previous systematic review of systematic reviews which may be eligible for inclusion for the map.

Elizabeth Ghogomu, Victoria Barbeau, Sierra Dowling, Rebecca Doyle, Ella Beveridge, Payaam Desai, Jimmy Huang, Nour Elmestekawy, Tarannum Hussain, Arpana Wadhvani, Sabrina

Boutin, Niobe Haitas, Douglas M. Salzwedel, Roger Simard, Paul Hebert, Christopher Mikton, have no conflicts of interest.

Plans for updating the EGM

The EGM will be updated every two years.

Differences between protocol and review

We simplified the conceptual framework in the review.

Characteristics of studies

Characteristics of included studies

See Supporting Information interactive EGM <https://globalageing.cochrane.org/cochrane-reviews>.

Characteristics of excluded studies

Studies	Reasons for exclusion
Bolle 2015 Clarkson 2018 Summer 2021	Evidence/aim (not to reduce social isolation or loneliness)
Dickens 2011 Gine-Garriga 2017 Jones 2019 Nicholson 2012 Toh 2016 Zeppegno 2018	Intervention (not digital)
Fan 2016* Forsman 2017* Gorenko 2021* Rebollar 2015 Perkins 2012 Preston 2019 Selak 2019* Winterton 2011*	Study design (not a systematic review of effectiveness* or a non-comparative primary study)
Burkow 2015 Cooper 2014 Dam 2017 Erfani 2018 Lara 2016 Nijman 2019	Target population (included < 60 years old, and not disaggregated by age)

--	--

References to studies

Included studies

Abbott R, Orr N, McGill P, Whear R, Bethel A, Garside R, Stein K, & Thompson-Coon J. (2019). How do “robopets” impact the health and well-being of residents in care homes? A systematic review of qualitative and quantitative evidence. *International Journal of Older People Nursing*, 14(3), e12239. <https://doi.org/10.1111/opn.12239>

Abdi J, Al-Hindawi A, Ng T, & Vizcaychipi M P. (2018). Scoping review on the use of socially assistive robot technology in elderly care. *BMJ Open*, 8(2), e018815. <https://doi.org/10.1136/bmjopen-2017-018815>

Abdi J, Al-Hindawi A, & Vizcaychipi M. (2017). Use of socially assistive robot technology in elderly care. *Age and Ageing*, 46 (Supplement 1), i1.

Abou Allaban, A, Wang M Z, & Padir T. (2020). A Systematic Review of Robotics Research in Support of In-Home Care for Older Adults. *Information*, 11(2). <https://doi.org/10.3390/info11020075>

Aggarwal B, Xiong Q, & Schroeder-Butterfill E. (2020). Impact of the use of the internet on quality of life in older adults: Review of literature. *Primary Health Care Research & Development*, 21, e55. <https://doi.org/10.1017/S1463423620000584>

Ali R, Hoque E, Duberstein P, Schubert L, Razavi S Z, Kane B, Silva C, Daks J S, Huang M, & Van Orden K. (2020). Aging and Engaging: A Pilot Randomized Controlled Trial of an Online Conversational Skills Coach for Older Adults. *American Journal of Geriatric Psychiatry*, 22, 22. <https://doi.org/10.1016/j.jagp.2020.11.004>

Alves G S, Bruno Costa, E, Alves G S, Rodrigues V M, Dourado M C. N, Casali M E, Carrilho C G, Veras A B, Rodrigues V M, & Dourado M C. N. (2020). A Systematic Review of Home-Setting Psychoeducation Interventions for Behavioral Changes in Dementia: Some

- Lessons for the COVID-19 Pandemic and Post-Pandemic Assistance. *Front. Psychiatry*, 11, 577871. <https://doi.org/10.3389/FPSYT.2020.577871>
- Archer N, Aria R, Assoc Comp, & Machinery. (2019). Reducing Risk from Chronic Illness with Life Style Changes Supported by Online Health Self-Management (p. 76). <https://doi.org/10.1109/I.2019.00020>
- Baez M, Khaghani Far, I, Ibarra F, Ferron M, Didino D, & Casati F. (2017). Effects of online group exercises for older adults on physical, psychological and social wellbeing: A randomized pilot trial. *PeerJ*, 5, e3150. <https://doi.org/10.7717/peerj.3150>
- Baker S, Warburton J, Waycott J, Batchelor F, Hoang T, Dow B, Ozanne E, & Vetere F. (2018). Combatting social isolation and increasing social participation of older adults through the use of technology: A systematic review of existing evidence. *Australasian Journal on Ageing*, 37(3), 184–193. <https://doi.org/10.1111/ajag.12572>
- Ballesteros Soledad, Toril Pilar, Mayas Julia, Reales Jose M, & Waterworth John A. (2014). An ICT-mediated social network in support of successful ageing. *Gerontechnology*, 13(1), 39–48. <https://doi.org/10.4017/gt.2014.13.1.007.00>
- Banks Marian R, Willoughby Lisa M, & Banks William A. (2008). Animal-assisted therapy and loneliness in nursing homes: Use of robotic versus living dogs. *Journal of the American Medical Directors Association*, 9(3), 173–177.
- Bedaf S, Gelderblom G J, & De Witte L. (2015). Overview and Categorization of Robots Supporting Independent Living of Elderly People: What Activities Do They Support and How Far Have They Developed. *Assistive Technology*, 27(2), 88–100. <https://doi.org/10.1080/10400435.2014.978916>
- Beentjes K M, Neal D P, Kerkhof Y J. F, Broeder C, Moeridjan Z D. J, Ettema T P, Pelkmans W, Muller M M, Graff M J. L, & Droes R M. (2020). Impact of the FindMyApps program on people with mild cognitive impairment or dementia and their caregivers; an exploratory pilot randomised controlled trial. *Disability & Rehabilitation Assistive Technology*, 1–13. <https://doi.org/10.1080/17483107.2020.1842918>

Bemelmans R, Gelderblom G J, Jonker P, & de Witte L. (2012). Socially assistive robots in elderly care: A systematic review into effects and effectiveness. *Journal of the American Medical Directors Association*, 13(2), 114-120.e1.

<https://doi.org/10.1016/j.jamda.2010.10.002>

Beneito-Montagut Roser, Cassián-Yde Nizaiá, & Begueria Arantza. (2018). What do we know about the relationship between internet-mediated interaction and social isolation and loneliness in later life? *Quality in Ageing and Older Adults*, 19(1), 14-30.

Bermeja Ana Isabel & Ausín Berta. (2018). Programas para combatir la soledad en las personas mayores en el ámbito institucionalizado: Una revisión de la literatura científica. *Revista Española de Geriátria y Gerontología*, 53(3), 155-164.

<https://doi.org/10.1016/j.regg.2017.05.006>

Bethell Jennifer, Babineau Jessica, Iaboni Andrea, Green Robin, Cuaresma-Canlas Rachelle, Karunanathan Roobika, Schon Barbara, Schon Denise, & McGilton Katherine S. (2019). Social integration and loneliness among long-term care home residents: Protocol for a scoping review. *BMJ Open*, 9(12). <https://doi.org/10.1136/bmjopen-2019-033240>

Boekhout J M, Volders E, Bolman C A. W, de Groot R H. M, & Lechner L. (2021). Long-Term Effects on Loneliness of a Computer-Tailored Intervention for Older Adults With Chronic Diseases: A Randomized Controlled Trial. *Journal of Aging & Health*, 8982643211015027. <https://doi.org/10.1177/08982643211015027>

Bornemann R. (2014). The impact of information and communication technology (ICT) usage on social isolation including loneliness in older adults. A systematic review. In Magdeburg/Stendal University of Applied Sciences.

<http://www.epistemonikos.org/documents/658a15ff5523ae6eb43a330d618cb3bfo89fe571>

Boston University. (2020). Impact of Covid-19 on Frequent Social Interaction Through Communication Technologies in the Cognitive Status of Socially-isolated Older Adults. *Clinicaltrials.Gov*. <https://clinicaltrials.gov/ct2/show/NCT04480112>

- Brodbeck J, Berger T, Biesold N, Rockstroh F, & Znoj H J. (2019). Evaluation of a guided internet-based self-help intervention for older adults after spousal bereavement or separation/divorce: A randomised controlled trial. *Journal of Affective Disorders*, 252, 440–449. <https://doi.org/10.1016/j.jad.2019.04.008>
- Bruce Martha L, Pepin Renee, Marti C Nathan, Stevens Courtney J, & Choi Namkee G. (2020). One Year Impact on Social Connectedness for Homebound Older Adults: Randomized Controlled Trial of Tele-delivered Behavioral Activation Versus Tele-delivered Friendly Visits. 28(7), 698–708. <https://doi.org/10.1016/j.jagp.2020.02.008>
- Buitenweg J I. V, Van De Ven R M, Ridderinkhof K R, & Murre J M. J. (2019). Does cognitive flexibility training enhance subjective mental functioning in healthy older adults? *Aging Neuropsychology & Cognition*, 26(5), 688–710. <https://doi.org/10.1080/13825585.2018.1519106>
- Cacciata M, Stromberg A, Lee J A, Sorkin D, Lombardo D, Clancy S, Nyamathi A, & Evangelista L S. (2019). Effect of exergaming on health-related quality of life in older adults: A systematic review. *International Journal of Nursing Studies*, 93, 30–40. <https://doi.org/10.1016/j.ijnurstu.2019.01.010>
- Campos W, Martinez A, Sanchez W, Estrada H, Castro-Sanchez N A, & Mujica D. (2016). A Systematic Review of Proposals for the Social Integration of Elderly People Using Ambient Intelligence and Social Networking Sites. *Cognitive Computation*, 8(3), 529–542. <https://doi.org/10.1007/s12559-016-9382-z>
- Casanova G, Zaccaria D, Rolandi E, & Guaita A. (2021). The Effect of Information and Communication Technology and Social Networking Site Use on Older People’s Well-Being in Relation to Loneliness: Review of Experimental Studies. *Journal of Medical Internet Research*, 23(3), e23588. <https://doi.org/10.2196/23588>
- Cattan Mima, White Martin, Bond John, & Learmouth Alison. (2005). Preventing social isolation and loneliness among older people: A systematic review of health promotion

- interventions. *Ageing and Society*, 25(1), 41–67.
<https://doi.org/10.1017/S0144686X04002594>
- Chang S M & Sung H C. C. (2013). The effectiveness of seal-like robot therapy on mood and social interactions of older adults: A systematic review protocol. *JBIC Library of Systematic Reviews*, 11(10), 68–75.
- Chen S C, Jones C, & Moyle W. (2018). Social Robots for Depression in Older Adults: A Systematic Review. *Journal of Nursing Scholarship*, 50(6), 612–622.
<https://doi.org/10.1111/jnu.12423>
- Chen Y R & Schulz P J. (2016). The Effect of Information Communication Technology Interventions on Reducing Social Isolation in the Elderly: A Systematic Review. *Journal of Medical Internet Research*, 18(1), e18. <https://doi.org/10.2196/jmir.4596>
- Chippis J, Jarvis M A, & Ramlall S. (2017). The effectiveness of e-Interventions on reducing social isolation in older persons: A systematic review of systematic reviews. *Journal of Telemedicine & Telecare*, 23(10), 817–827. <https://doi.org/10.1177/1357633X17733773>
- Chiu C J & Wu C H. (2019). Information and Communications Technology as a Health Promotion Method for Older Adults in Assisted-Living Facilities: Three-Arm Group-Randomized Trial. 24058297, 2(1), e12633. <https://doi.org/10.2196/12633>
- Choi H K & Lee S H. (2021). Trends and Effectiveness of ICT Interventions for the Elderly to Reduce Loneliness: A Systematic Review. *Healthcare*, 9(3), 07.
<https://doi.org/10.3390/healthcare9030293>
- Choi M, Kong S, & Jung D. (2012). Computer and internet interventions for loneliness and depression in older adults: A meta-analysis. *Healthcare Informatics Research*, 18(3), 191–198. <https://doi.org/10.4258/hir.2012.18.3.191>
- Choi N G, Marti C N, Wilson N L, Chen G J, Sirrianni L, Hegel M T, Bruce M L, & Kunik M E. (2020). Effect of Telehealth Treatment by Lay Counselors vs by Clinicians on Depressive Symptoms Among Older Adults Who Are Homebound: A Randomized Clinical Trial.

JAMA Network Open, 3(8), 2574–3805.

<https://doi.org/10.1001/jamanetworkopen.2020.15648>

Choi N G, Pepin R, Marti C N, Stevens C J, & Bruce M L. (2020). Improving Social Connectedness for Homebound Older Adults: Randomized Controlled Trial of Tele-Delivered Behavioral Activation Versus Tele-Delivered Friendly Visits. *American Journal of Geriatric Psychiatry*, 28(7), 698–708. <https://doi.org/10.1016/j.jagp.2020.02.008>

Chua Puay-Hoe, Jung Younbo, Lwin May O, & Theng Yin-Leng. (2013). Let's play together: Effects of video-game play on intergenerational perceptions among youth and elderly participants. *Computers in Human Behavior*, 29(6), 2303–2311.

Cohen-Mansfield Jiska & Perach Rotem. (2015). Interventions for alleviating loneliness among older persons: A critical review. 29(3), e109-25. <https://doi.org/10.4278/ajhp.130418-lit-182>

Coll-Planas L, Nyqvist F, Puig T, Urrútia G, Solà I, & Montserrat R. (2017). Social capital interventions targeting older people and their impact on health: A systematic review. *Journal of Epidemiology and Community Health*, 71(7), 663–672. <https://doi.org/10.1136/jech-2016-208131>

Courtin Emilie & Knapp Martin. (2015). Social isolation, loneliness and health in old age: A scoping review. 25(3), 799–812. <https://doi.org/10.1111/hsc.12311>

Czaja S J, Boot W R, Charness N, Rogers W A, & Sharit J. (2018). Improving Social Support for Older Adults Through Technology: Findings From the PRISM Randomized Controlled Trial. *Gerontologist*, 58(3), 467–477. <https://doi.org/10.1093/geront/gnw249>

Czaja S J, Boot W R, Charness N, W A Rogers, Sharit J, Fisk A D, Lee C C, & Nair S N. (2015). The personalized reminder information and social management system (PRISM) trial: Rationale, methods and baseline characteristics. *Contemporary Clinical Trials*, 40, 35–46. <https://doi.org/10.1016/j.cct.2014.11.004>

- Da-Eun K I. M, Hyang K I. M, Junghee Hyun, Hyojin L E. E, Hye Hyun Sung, Soyoung B A. E, Sunghee H Tak, Yeon-Hwan Park, & Ju-Young Yoon. (2018). Interventions Using Technologies for Older Adults in Long-term Care Facilities: A Systematic Review. *Journal of Korean Academy of Community Health Nursing*, 170–183.
- Damant Jacqueline, Knapp Martin, Freddolino Paul, & Lombard Daniel. (2017). Effects of digital engagement on the quality of life of older people. *Health & Social Care in the Community*, 25(6), 1679–1703. <https://doi.org/10.1111/hsc.12335>
- D’Cunha N M, Isbel S T, Frost J, Fearon A, McKune A J, Naumovski N, & Kellett J. (n.d.). Effects of a virtual group cycling experience on people living with dementia: A mixed method pilot study. *Dementia-International Journal of Social Research and Practice*. <https://doi.org/10.1177/1471301220951328>
- Dequanter S, Gagnon M P, Ndiaye M A, Gorus E, Fobelets M, Giguere A, Bourbonnais A, & Buyl R. (2020). The Effectiveness of E-Health Solutions for Ageing With Cognitive Impairment: A Systematic Review. *Gerontologist*, 11, 11. <https://doi.org/10.1093/geront/gnaa065>
- Dermody G, Whitehead L, Wilson G, & Glass C. (2020). The Role of Virtual Reality in Improving Health Outcomes for Community-Dwelling Older Adults: Systematic Review. *Journal of Medical Internet Research*, 22(6), e17331. <https://doi.org/10.2196/17331>
- Dichter M N, Albers B, Trutschel D, Strobel A M, Seismann-Petersen S, Wermke K, Halek M, & Berwig M. (2020). TALKING TIME: A pilot randomized controlled trial investigating social support for informal caregivers via the telephone. *BMC Health Services Research*, 20(1), 788. <https://doi.org/10.1186/s12913-020-05523-9>
- Dickens A P, Richards S H, Greaves C J, & Campbell J L. (2011). Interventions targeting social isolation in older people: A systematic review. *BMC Public Health*, 11, 647. <https://doi.org/10.1186/1471-2458-11-647>

Dodge H, Goodrich E, Lindsley J, Hampstead B, & Kaye J. (2018). Behavioral RCT using frequent social interaction to enhance cognitive reserve. *Clinical Trials*, 15 (Supplement 2), 130–131.

Dodge H H, Zhu J, Mattek N, Bowman M, Ybarra O, Wild K, Loewenstein D A, & Kaye J A. (2015). Web-enabled Conversational Interactions as a Means to Improve Cognitive Functions: Results of a 6-Week Randomized Controlled Trial. *Alzheimer's & Dementia : Translational Research & Clinical Interventions*, 1(1), 1–12.
<https://doi.org/10.1016/j.trci.2015.01.001>

Duggleby W, O'Rourke H, Swindle J, Peacock S, McAiney C, Baxter P, Thompson G, Dube V, Nekolaichuk C, Ghosh S, & Holroyd-Leduc J. (2020). Study protocol: Pragmatic randomized control trial of my tools 4 care- in care (MT4C-in care) a web-based tool for family Carers of persons with dementia residing in long term care. *BMC Geriatrics*, 20(1), 285. <https://doi.org/10.1186/s12877-020-01690-w>

El Kamali M, Angelini L, Caon M, Carrino F, Rocke C, Guye S, Rizzo G, Mastropietro A, Sykora M, Elayan S, Kniestedt I, Ziylan C, Lettieri E, Abou Khaled, O, & Mugellini E. (2020). Virtual Coaches for Older Adults' Wellbeing: A Systematic Review. *Ieee Access*, 8, 101884–101902. <https://doi.org/10.1109/access.2020.2996404>

Engelbrecht Romy & Shoemark Helen. (2015). The Acceptability and Efficacy of Using iPads in Music Therapy to Support Wellbeing with Older Adults: A Pilot Study. *The Australian Journal of Music Therapy*, 26, 49–70.

Etxeberria I, Salaberria K, & Gorostiaga A. (n.d.). Online support for family caregivers of people with dementia: A systematic review and meta-analysis of RCTs and quasi-experimental studies. *Aging & Mental Health*. <https://doi.org/10.1080/13607863.2020.1758900>

Fakoya Olujoke A, McCorry Noleen K, & Donnelly Michael. (2020). Loneliness and social isolation interventions for older adults: A scoping review of reviews. 20(1), 129–129.
<https://doi.org/10.1186/s12889-020-8251-6>

- Feng Y, Barakova E I, Yu S, Hu J, & Rauterberg G W. M. (2020). Effects of the Level of Interactivity of a Social Robot and the Response of the Augmented Reality Display in Contextual Interactions of People with Dementia. *Sensors*, 20(13), 05. <https://doi.org/10.3390/s20133771>
- Ferreira Sónia, Torres Ana, Mealha Óscar, & Veloso Ana. (2015). Training Effects on Older Adults in Information and Communication Technologies Considering Psychosocial Variables. *Educational Gerontology*, 41(7), 482–493. <https://doi.org/10.1080/03601277.2014.994351>
- Fields J, Cembali A G, Michalec C, Uchida D, Griffiths K, Cardes H, Cuellar J, Chodos A H, & Lyles C R. (2021). In-Home Technology Training Among Socially Isolated Older Adults: Findings From the Tech Allies Program. *Journal of Applied Gerontology*, 40(5), 489–499. <https://doi.org/10.1177/0733464820910028>
- Fokkema T & Knipscheer K. (2007). Escape loneliness by going digital: A quantitative and qualitative evaluation of a Dutch experiment in using ECT to overcome loneliness among older adults. *Aging & Mental Health*, 11(5), 496–504. <https://doi.org/10.1080/13607860701366129>
- Forsman A K, Nordmyr J, Matosevic T, Park A L, Wahlbeck K, & McDaid D. (2018). Promoting mental wellbeing among older people: Technology-based interventions. *Health Promotion International*, 33(6), 1042–1054. <https://doi.org/10.1093/heapro/dax047>
- Francis J, Rikard R V, Cotten S R, & Kadylak T. (2019). Does ICT Use matter? How information and communication technology use affects perceived mattering among a predominantly female sample of older adults residing in retirement communities. *Information Communication & Society*, 22(9), 1281–1294. <https://doi.org/10.1080/1369118x.2017.1417459>
- Gardiner Clare, Geldenhuys Gideon, & Gott Meryn. (2018). Interventions to reduce social isolation and loneliness among older people: An integrative review. *Health & Social Care in the Community*, 26(2), 147–157. <https://doi.org/10.1111/hsc.12367>

- Godwin K M, Mills W L, Anderson J A, & Kunik M E. (2013). Technology-driven interventions for caregivers of persons with dementia: A systematic review. *American Journal of Alzheimer's Disease & Other Dementias*, 28(3), 216–222. <https://doi.org/10.1177/1533317513481091>
- Gongora Alonso, S, Hamrioui S, de la Torre Diez I, Motta Cruz, E, Lopez-Coronado M, & Franco M. (2019). Social Robots for People with Aging and Dementia: A Systematic Review of Literature. *Telemedicine Journal & E-Health*, 25(7), 533–540. <https://doi.org/10.1089/tmj.2018.0051>
- Gustafson D H, Gustafson D H, Cody O J, Chih M Y, Johnston D C, & Asthana S. (2019). Pilot Test of a Computer-Based System to Help Family Caregivers of Dementia Patients. *Journal of Alzheimer's Disease*, 70(2), 541–552. <https://doi.org/10.3233/JAD-190052>
- Gustafson D H, Sr, Mares M L, Johnston D C, Mahoney J E, Brown R T, Landucci G, Pe-Romashko K, Cody O J, Gustafson D H, Jr, & Shah D V. (2021). A Web-Based eHealth Intervention to Improve the Quality of Life of Older Adults With Multiple Chronic Conditions: Protocol for a Randomized Controlled Trial. *JMIR Research Protocols*, 10(2), e25175. <https://doi.org/10.2196/25175>
- Gustafson D H, Sr, McTavish F, Gustafson D H, Jr, Mahoney J E, Johnson R A, Lee J D, Quanbeck A, Atwood A K, Isham A, Veeramani R, Clemson L, & Shah D. (2021). Effect of an eHealth intervention on older adults' quality of life and health-related outcomes: A randomized clinical trial. *Journal of General Internal Medicine*, 1–10. <https://doi.org/10.1007/s11606-021-06888-1>
- Haase KR, Sattar S, Hall S, McLean B, Wills A, Gray M, Kenis C, Donison V, Howell D, & Puts M. (2021). Systematic review of self-management interventions for older adults with cancer. *Psycho-Oncology*, 30(7), 989–1008. <https://doi.org/10.1002/pon.5649>
- Hagan R, Manktelow R, Taylor B J, & Mallett J. (2014). Reducing loneliness amongst older people: A systematic search and narrative review. *Aging & Mental Health*, 18(6), 683–693. <https://doi.org/10.1080/13607863.2013.875122>

- Hartke R J & King R B. (2003). Telephone group intervention for older stroke caregivers. *Topics in Stroke Rehabilitation*, 9(4), 65–81.
- Health Quality & Ontario. (2008). Social isolation in community-dwelling seniors: An evidence-based analysis. *Ontario Health Technology Assessment Series*, 8(5), 1–49.
- Heller K, Thompson M G, Trueba P E, Hogg J R, & Vlachos-Weber I. (1991). Peer support telephone dyads for elderly women: Was this the wrong intervention? *American Journal of Community Psychology*, 19(1), 53–74.
- Hernández-Ascanio José, Pérula-de Torres Luis Ángel, Roldán-Villalobos Ana, Pérula-de Torres José Carlos, & Rich-Ruiz Manuel. (2020). Effectiveness of a multicomponent intervention to reduce social isolation and loneliness in community-dwelling elders: A randomized clinical trial. Study protocol. *Journal of Advanced Nursing*, 76(1), 337–346. <https://doi.org/10.1111/jan.14230>
- Heyn Billipp & S. (2001). The psychosocial impact of interactive computer use within a vulnerable elderly population: A report on a randomized prospective trial in a home health care setting. *Public Health Nursing*, 18(2), 138–145.
- Hicken B L, Daniel C, Luptak M, Grant M, Kilian S, & Rupper R W. (2017). Supporting Caregivers of Rural Veterans Electronically (SCORE). *Journal of Rural Health*, 33(3), 305–313. <https://doi.org/10.1111/jrh.12195>
- Hoang P, Whaley C, Thompson K, Ho V, Rehman U, Boluk K, & Grindrod K A. (2021). Evaluation of an Intergenerational and Technological Intervention for Loneliness: Protocol for a Feasibility Randomized Controlled Trial. *JMIR Research Protocols*, 10(2), e23767. <https://doi.org/10.2196/23767>
- Hopwood Jenny, Walker Nina, McDonagh Lorraine, Rait Greta, Walters Kate, Iliffe Stephen, Ross Jamie, & Davies Nathan. (2018). Internet-based interventions aimed at supporting family caregivers of people with dementia: Systematic review. *Journal of Medical Internet Research* Vol 20(6), 2018, ArtID E216, 20(6). <https://www.jmir.org/2018/6/e216/>

- Hung L, Chow B, Shadarevian J, O'Neill R, Berndt A, Wallsworth C, Horne N, Gregorio M, Mann J, Son C, & Chaudhury H. (2021). Using touchscreen tablets to support social connections and reduce responsive behaviours among people with dementia in care settings: A scoping review. *Dementia*, 20(3), 1124–1143.
- Hung Lillian, Liu Cindy, Woldum Evan, Au-Yeung Andy, Berndt Annette, Wallsworth Christine, Horne Neil, Gregorio Mario, Mann Jim, & Chaudhury Habib. (2019). The benefits of and barriers to using a social robot PARO in care settings: A scoping review. 19(1), 232–232. <https://doi.org/10.1186/s12877-019-1244-6>
- Ibarra F, Baez M, Cernuzzi L, & Casati F. (2020). A Systematic Review on Technology-Supported Interventions to Improve Old-Age Social Wellbeing: Loneliness, Social Isolation, and Connectedness. *Journal of Healthcare Engineering*, 2020, 2036842. <https://doi.org/10.1155/2020/2036842>
- Ibrahim A, Chong M C, Khoo S, Wong L P, Chung I, & Tan M P. (2021). Virtual Group Exercises and Psychological Status among Community-Dwelling Older Adults during the COVID-19 Pandemic-A Feasibility Study. *Geriatrics*, 6(1), 22. <https://doi.org/10.3390/geriatrics6010031>
- Iman Khaghani, Far, Ferron Michela, Ibarra Francisco, Baez Marcos, Tranquillini Stefano, Casati Fabio, & Doppio Nicola. (2015). The interplay of physical and social wellbeing in older adults: Investigating the relationship between physical training and social interactions with virtual social environments. *PeerJ Computer Science*. <https://doi.org/10.7717/peerj-cs.30>
- Isabet B, Pino M, Lewis M, Benveniste S, & Rigaud A S. (2021). Social Telepresence Robots: A Narrative Review of Experiments Involving Older Adults before and during the COVID-19 Pandemic. *International Journal of Environmental Research & Public Health* [Electronic Resource], 18(7), 30. <https://doi.org/10.3390/ijerph18073597>
- Jackson David, Roberts Gail, Wu Min Lin, Ford Rosemary, & Doyle Colleen. (2016). A systematic review of the effect of telephone, internet or combined support for carers of

people living with Alzheimer's, vascular or mixed dementia in the community. *Archives of Gerontology and Geriatrics*, 66, 218–236.

- Jarvis M A, Padmanabhanunni A, & Chipps J. (2019). An Evaluation of a Low-Intensity Cognitive Behavioral Therapy mHealth-Supported Intervention to Reduce Loneliness in Older People. *International Journal of Environmental Research & Public Health* [Electronic Resource], 16(7), 11. <https://doi.org/10.3390/ijerph16071305>
- Jarvis Mary Ann, Padmanabhanunni Anita, Balakrishna Yusenantha, & Chipps Jennifer. (2020). The effectiveness of interventions addressing loneliness in older persons: An umbrella review. 12(NA), 100177-NA. <https://doi.org/10.1016/j.ijans.2019.100177>
- Jones C, Liu D F, Murfield M J, & Moyle P W. (2020). Effects of non-facilitated meaningful activities for people with dementia in long-term care facilities: A systematic review. *Geriatric Nursing* (New York, N.Y.). <https://doi.org/10.1016/j.gerinurse.2020.06.001>
- Jones R B, Ashurst E J, Atkey J, & Duffy B. (2015). Older people going online: Its value and before-after evaluation of volunteer support. *Journal of Medical Internet Research*, 17(5), e122. <https://doi.org/10.2196/jmir.3943>
- Kachouie R, Sedighadeli S, & Abkenar A B. (2017). The Role of Socially Assistive Robots in Elderly Wellbeing: A Systematic Review. In *Cross-Cultural Design* (Vol. 10281, pp. 669–682). https://doi.org/10.1007/978-3-319-57931-3_54
- Kachouie R, Sedighadeli S, Khosla R, & Chu M T. (2014). Socially Assistive Robots in Elderly Care: A Mixed-Method Systematic Literature Review. *International Journal of Human-Computer Interaction*, 30(5), 369–393. <https://doi.org/10.1080/10447318.2013.873278>
- Kahlon M K, Aksan N, Aubrey R, Clark N, Cowley-Morillo M, Jacobs E A, Mundhenk R, Sebastian K R, & Tomlinson S. (2021). Effect of Layperson-Delivered, Empathy-Focused Program of Telephone Calls on Loneliness, Depression, and Anxiety among Adults during the COVID-19 Pandemic: A Randomized Clinical Trial. *JAMA Psychiatry*. <https://jamanetwork.com/journals/jamapsychiatry/fullarticle/2776786>

- Kamalpour M, Watson J, & Buys L. (2020). How Can Online Communities Support Resilience Factors among Older Adults. *International Journal of Human-Computer Interaction*, 36(14), 1342–1353. <https://doi.org/10.1080/10447318.2020.1749817>
- kazazi Leila, Shati Mohsen, Mortazavi Seyede Salehe, Nejati Vahid, & Foroughan Mahshid. (2021). The impact of computer-based cognitive training intervention on the quality of life among elderly people: A randomized clinical trial. *Trials*, 22(1), 1–10. <https://doi.org/10.1186/s13063-020-05008-4>
- Keogh Justin W, Power Nicola, Wooller Leslie, Lucas Patricia, & Whatman Chris. (2014). Physical and psychosocial function in residential aged-care elders: Effect of Nintendo Wii sports games. *Journal of Aging and Physical Activity*, 22(2), 235–244.
- Khosravi P & Ghananchi A H. (2016). Investigating the effectiveness of technologies applied to assist seniors: A systematic literature review. *International Journal of Medical Informatics*, 85(1), 17–26. <https://doi.org/10.1016/j.ijmedinf.2015.05.014>
- Khosravi Pouria, Rezvani Azadeh, & Wiewiora Anna. (2016). The impact of technology on older adults' social isolation. *Computers in Human Behavior*, 63, 594–603. <https://doi.org/10.1016/j.chb.2016.05.092>
- Koh W Q, Felding S A, Toomey E, & Casey D. (2021). Barriers and facilitators to the implementation of social robots for older adults and people with dementia: A scoping review protocol. *Systematic Reviews*, 10(1), 49. <https://doi.org/10.1186/s13643-021-01598-5>
- Salehi A, Salehi E, Mosadeghi-Nik M, Sargeant S, & Fatehi F. (n.d.). Strengthening positive social pathways via digital social applications in individuals with social skills deficits: A scoping review. *International Journal of Social Psychiatry*. <https://doi.org/10.1177/0020764020963354>
- Koh W Q, Ang F X. H, & Casey D. (2021). Impacts of Low-cost Robotic Pets for Older Adults and People With Dementia: Scoping Review. *JMIR Rehabilitation And Assistive Technologies*, 8(1), e25340. <https://doi.org/10.2196/25340>

- Koo B M & Vizer L M. (2019). Examining Mobile Technologies to Support Older Adults With Dementia Through the Lens of Personhood and Human Needs: Scoping Review. *JMIR MHealth and UHealth*, 7(11), e15122. <https://doi.org/10.2196/15122>
- Kramer L L, Mulder B C, van Velsen L, & de Vet E. (2021). Use and Effect of Web-Based Embodied Conversational Agents for Improving Eating Behavior and Decreasing Loneliness Among Community-Dwelling Older Adults: Protocol for a Randomized Controlled Trial. *JMIR Research Protocols*, 10(1), e22186. <https://doi.org/10.2196/22186>
- Kubra Koce & MSc P T. (2020). Telerehabilitation in Individuals Over 65 Years of Age Having Social Isolation Due to Coronavirus (Covid-19). *Clinicaltrials.Gov*. <https://clinicaltrials.gov/ct2/show/NCT04334434?term=NCT04334434&draw=2&rank=1>
- Lady Davis & Institute. (2021). Virtual Chair Yoga for Older Adults and Caregivers: Randomized Controlled Trial. *Clinicaltrials.Gov*. <https://clinicaltrials.gov/ct2/show/NCT04884529>
- Laganà Luciana & García James J. (2013). The Mental Health Impact of Computer and Internet Training on a Multi-ethnic Sample of Community-Dwelling Older Adults: Results of a Pilot Randomised Controlled Trial. 9(3), 135–147.
- Lai Daniel W. L, Li Jia, Ou Xiaoting, & Li Celia Y. P. (2020). Effectiveness of a peer-based intervention on loneliness and social isolation of older Chinese immigrants in Canada: A randomized controlled trial. *BMC Geriatrics*, 20, 1–12. <https://doi.org/10.1186/s12877-020-01756-9>
- Lappalainen P, Pakkala I, Lappalainen R, & Nikander R. (2021). Supported Web-Based Acceptance and Commitment Therapy for Older Family Caregivers (CareACT) Compared to Usual Care. *Clinical Gerontologist*, 1–17. <https://doi.org/10.1080/07317115.2021.1912239>

Larsson Ellinor, Padyab Mojgan, Larsson-Lund Maria, & Nilsson Ingeborg. (2016). Effects of a social internet-based intervention programme for older adults: An explorative randomised crossover study. *British Journal of Occupational Therapy*, 79(10), 629–636. <https://doi.org/10.1177/0308022616641701>

Lauriks S, Reinersmann A, van der Roest H G, Meiland Franka, Davies Richard, Moelaert F, Mulvenna Maurice, Nugent Chris D, & Dröes Rose-Marie. (2007). Review of ICT-Based Services for Identified Unmet Needs in People with Dementia. 6(3), 223–246. <https://doi.org/10.1016/j.arr.2007.07.002>

Lawson Health Research & Institute. (2021). SSIPP vs. PST vs. WLC. *Clinicaltrials.Gov*. <https://clinicaltrials.gov/ct2/show/NCT04887350>

Lee C, Kuhn I, McGrath M, Remes O, Cowan A, Duncan F, Baskin C, Oliver E J, Osborn D P. J, Dykxhoorn J, Kaner E, Walters K, Kirkbride J, Gnani S, Lafortune L, & Programme Nihhr Sphr Public Mental Health. (2021). A systematic scoping review of community-based interventions for the prevention of mental ill-health and the promotion of mental health in older adults in the UK. *Health & Social Care in the Community*, 14, 14. <https://doi.org/10.1111/hsc.13413>

Li J, Xu X, Pham T P, Theng Y L, Katajapuu N, & Luimula M. (2017). Exergames Designed for Older Adults: A Pilot Evaluation on Psychosocial Well-Being. *Games for Health Journal*, 6(6), 371–378. <https://doi.org/10.1089/g4h.2017.0072>

Li Jinhui, Erdt Mojisola, Chen Luxi, Cao Yuanyuan, Lee Shan-Qi, & Theng Yin-Leng. (2018). The social effects of exergames on older adults: Systematic review and metric analysis. *Journal of Medical Internet Research* Vol 20(6), 2018, ArtID E10486, 20(6). <https://login.proxy.bib.uottawa.ca/login?url=http://ovidsp.ovid.com/ovidweb.cgi?T=JS&CSC=Y&NEWS=N&PAGE=fulltext&D=psyc15&AN=2018-57129-001>

Liang A, Piroth I, Robinson H, MacDonald B, Fisher M, Nater U M, Skoluda N, & Broadbent E. (2017). A Pilot Randomized Trial of a Companion Robot for People With Dementia Living in the Community. *Journal of the American Medical Directors Association*, 18(10), 871–878. <https://doi.org/10.1016/j.jamda.2017.05.019>

- Lin T Y, Huang C M, Hsu H P, Liao J Y, Cheng V Y, Wang S W, & Guo J L. (2020). Effects of a Combination of Three-Dimensional Virtual Reality and Hands-on Horticultural Therapy on Institutionalized Older Adults' Physical and Mental Health: Quasi-Experimental Design. *Journal of Medical Internet Research*, 22(11), e19002.
<https://doi.org/10.2196/19002>
- Lins S, Hayder-Beichel D, Rucker G, Motschall E, Antes G, Meyer G, & Langer G. (2014). Efficacy and experiences of telephone counselling for informal carers of people with dementia. *Cochrane Database of Systematic Reviews*, 9, CD009126.
<https://doi.org/10.1002/14651858.CD009126.pub2>
- Lopez-Hartmann M, Wens J, Verhoeven V, & Remmen R. (2012). The effect of caregiver support interventions for informal caregivers of community-dwelling frail elderly: A systematic review. *International Journal of Integrated Care*, 12, e133.
<https://doi.org/10.5334/ijic.845>
- Masi Christopher M, Chen Hsi-Yuan, Hawkey Louise C, & Cacioppo John T. (2010). A Meta-Analysis of Interventions to Reduce Loneliness: 15(3), 219–266.
<https://doi.org/10.1177/1088868310377394>
- Matson Theresa E, Anderson Melissa L, Renz Anne D, Greenwood-Hickman Mikael Anne, McClure Jennifer B, & Rosenberg Dori E. (2019). Changes in self-reported health and psychosocial outcomes in older adults enrolled in sedentary behavior intervention study. *American Journal of Health Promotion*, 33(7), 1053–1057.
<https://doi.org/10.1177/0890117119841405>
- Matz-Costa C, Lubben J, Lachman M E, Lee H, & Choi Y J. (2018). A Pilot Randomized Trial of an Intervention to Enhance the Health-Promoting Effects of Older Adults' Activity Portfolios: The Engaged4Life Program. *Journal of Gerontological Social Work*, 61(8), 792–816. <https://doi.org/10.1080/01634372.2018.1542371>

- McKechnie V, Barker C, & Stott J. (2014). Effectiveness of computer-mediated interventions for informal carers of people with dementia-a systematic review. *International Psychogeriatrics*, 26(10), 1619–1637. <https://doi.org/10.1017/s1041610214001045>
- Mikkelsen A S. B, Petersen S, Dragsted A C, & Kristiansen M. (2019). Social Interventions Targeting Social Relations Among Older People at Nursing Homes: A Qualitative Synthesized Systematic Review. *Inquiry*, 56, 46958018823929. <https://doi.org/10.1177/0046958018823929>
- Milbury K, Higgins H M, Chen A B, Antonoff M, Li Y, Tsao A S, & Bruera E. (2020). Pilot randomized controlled trial (RCT) in women with non-small cell lung cancer (NSCLC) to assess the feasibility of delivering group-based psychosocial care via videoconference. *Journal of Clinical Oncology. Conference*, 38(29). <https://www.liebertpub.com/doi/abs/10.1089/g4h.2017.0072>
- Miller K J, Adair B S, Pearce A J, Said C M, Ozanne E, & Morris M M. (2014). Effectiveness and feasibility of virtual reality and gaming system use at home by older adults for enabling physical activity to improve health-related domains: A systematic review. *Age & Ageing*, 43(2), 188–195. <https://doi.org/10.1093/ageing/aft194>
- Mittelman M S, Haley W E, Clay O J, & Roth D L. (2006). Improving caregiver well-being delays nursing home placement of patients with Alzheimer disease. *Neurology*, 67(9), 1592–1599. <https://doi.org/10.1212/01.wnl.0000242727.81172.91>
- Montana J I, Matamala-Gomez M, Maisto M, Mavrodiev P A, Cavalera C M, Diana B, Mantovani F, & Realdon O. (2020). The Benefits of emotion Regulation Interventions in Virtual Reality for the Improvement of Wellbeing in Adults and Older Adults: A Systematic Review. *Journal of Clinical Medicine*, 9(2), 12. <https://doi.org/10.3390/jcm9020500>
- Morris M E, Adair B, Ozanne E, Kurowski W, Miller K J, Pearce A J, Santamaria N, Long M, Ventura C, & Said C M. (2014). Smart technologies to enhance social connectedness in older people who live at home. *Australasian Journal on Ageing*, 33(3), 142–152. <https://doi.org/10.1111/ajag.12154>

- Morton T A, Wilson N, Haslam C, Birney M, Kingston R, & McCloskey L G. (2018). Activating and Guiding the Engagement of Seniors With Online Social Networking: Experimental Findings From the AGES 2.0 Project. *Journal of Aging & Health*, 30(1), 27–51. <https://doi.org/10.1177/0898264316664440>
- Mountain G A, Hind D, Gossage-Worrall R, Walters S J, Duncan R, Newbould L, Rex S, Jones C, Bowling A, Cattan M, Cairns A, Cooper C, Edwards R T, & Goyder E C. (2014). “Putting Life in Years” (PLINY) telephone friendship groups research study: Pilot randomised controlled trial. *Trials [Electronic Resource]*, 15, 141. <https://doi.org/10.1186/1745-6215-15-141>
- Moyle W, Jones C, Murfield J, Thalib L, Beattie E, Shum D, O’Dwyer S, Mervin C, & Draper B. (2017). Use of a Robotic Seal as a Therapeutic Tool to Improve Dementia Symptoms: A Cluster-Randomized Controlled Trial. *Journal of the American Medical Directors Association*, 18(9), 766–773. <https://doi.org/10.1016/j.jamda.2017.03.018>
- Myhre J W, Mehl M R, & Glisky E L. (2017). Cognitive Benefits of Online Social Networking for Healthy Older Adults. *Journals of Gerontology Series B-Psychological Sciences & Social Sciences*, 72(5), 752–760. <https://doi.org/10.1093/geronb/gbw025>
- Myhre Janelle Wohltmann. (2015). Effects of online social networking on the cognitive, social, and emotional health of older adults. *Dissertation Abstracts International: Section B: The Sciences and Engineering*, 76(1-B(E)). <https://repository.arizona.edu/handle/10150/332671>
- Neal D P, Kerkhof Y J. F, Ettema T P, Muller M, Bosmans J, Finnema E, Graff M, Dijkstra K, Stek M L, & Drees R M. (2021). Evaluation of FindMyApps: Protocol for a randomized controlled trial of the effectiveness and cost-effectiveness of a tablet-based intervention to improve self-management and social participation of community-dwelling people with mild dementia, compared to usual tablet use. *BMC Geriatrics*, 21(1), 138. <https://doi.org/10.1186/s12877-021-02038-8>

- Nef T, Ganea R L, Muri R M, & Mosimann U P. (2013). Social networking sites and older users—A systematic review. *International Psychogeriatrics*, 25(7), 1041–1053.
<https://doi.org/10.1017/S1041610213000355>
- Nijhof N, van Gemert-Pijnen J, Dohmen D, & Seydel E. (2009). Dementia and technology: A study of technology interventions in the healthcare for dementia patients and their caregivers. *Tijdschrift Voor Gerontologie En Geriatrie*, 40(3), 113–132.
- Nikitina S, Didino D, Baez M, & Casati F. (2018). Feasibility of Virtual Tablet-Based Group Exercise Among Older Adults in Siberia: Findings From Two Pilot Trials. *JMIR MHealth and UHealth*, 6(2), e40. <https://doi.org/10.2196/mhealth.7531>
- Nilsson M Y, Andersson S, Magnusson L, & Hanson E. (2021). Ambient assisted living technology-mediated interventions for older people and their informal carers in the context of healthy ageing: A scoping review. *Health Science Reports*, 4(1), e225.
<https://doi.org/10.1002/hsr2.225>
- Noone C, McSharry J, Smalle M, Burns A, Dwan K, Devane D, & Morrissey E C. (2020). Video calls for reducing social isolation and loneliness in older people: A rapid review. *Cochrane Database of Systematic Reviews*, 5, CD013632.
<https://doi.org/10.1002/14651858.CD013632>
- Ollevier A, Aguiar G, Palomino M, & Simpelaere I S. (2020). How can technology support ageing in place in healthy older adults? A systematic review. *Public Health Reviews*, 41(1), 26.
<https://doi.org/10.1186/s40985-020-00143-4>
- O'Rourke H M, Collins L, & Sidani S. (2018). Interventions to address social connectedness and loneliness for older adults: A scoping review. *BMC Geriatrics*, 18(1), 214.
<https://doi.org/10.1186/s12877-018-0897-x>
- Papadopoulos C, Castro N, Nigath A, Davidson R, Faulkes N, Menicatti R, Khaliq A A, Recchiuto C, Battistuzzi L, Randhawa G, Merton L, Kanoria S, Chong N Y, Kamide H, Hewson D, & Sgorbissa A. (2021). The CARESSES Randomised Controlled Trial: Exploring the Health-Related Impact of Culturally Competent Artificial Intelligence Embedded Into

Socially Assistive Robots and Tested in Older Adult Care Homes. *International Journal of Social Robotics*, 1–12. <https://doi.org/10.1007/s12369-021-00781-x>

Parkinson L, Banbury A, Livingstone A, Gordon S, Ray B, Byrne L, Nancarrow S, Doran C, McAllister M, Petersen C, Pedell S, & Wood D. (2018). Caring for Carers of People with Dementia: A Protocol for Harnessing Innovation Through Deploying Leading Edge Technologies to Enable Virtual Support Groups and Services. *Studies in Health Technology & Informatics*, 246, 29–41.

Pedrozo Campos Antunes, T, Souza Bulle de Oliveira, A., Hudec R, Brusque Crocetta, T, Ferreira de Lima Antao, J. Y., de Almeida Barbosa R. T, Guarnieri R, Massetti T, Garner D M, & de Abreu L C. (2019). Assistive technology for communication of older adults: A systematic review. *Aging & Mental Health*, 23(4), 417–427. <https://doi.org/10.1080/13607863.2018.1426718>

Pepin R, DiMilia P, Choi N, & Bruce M. (2019). Tele-Behavioral Activation for Social Isolation in Older Home-Delivered Meals Recipients: Preliminary Results from an Ongoing Randomized Controlled Trial. *American Journal of Geriatric Psychiatry*, 27 (3 Supplement), S129–S130.

Pereira L, Dias A, Queiros A, & Rocha N P. (2018). Technologies for Ageing in Place: A Systematic Review of Reviews and Meta-analyses. In *Biomedical Engineering Systems and Technologies* (Vol. 881, pp. 331–353). https://doi.org/10.1007/978-3-319-94806-5_18

Peres K, Zamudio-Rodriguez A, Dartigues J F, Amieva H, & Lafitte S. (2021). Prospective pragmatic quasi-experimental study to assess the impact and effectiveness of an innovative large-scale public health intervention to foster healthy ageing in place: The SoBeezy program protocol. *BMJ Open*, 11(4), e043082. <https://doi.org/10.1136/bmjopen-2020-043082>

Pinto-Bruno A C, Garcia-Casal J A, Csipke E, Jenaro-Rio C, & Franco-Martin M. (2017). ICT-based applications to improve social health and social participation in older adults with

- dementia. A systematic literature review. *Aging & Mental Health*, 21(1), 58–65.
<https://doi.org/10.1080/13607863.2016.1262818>
- Portz J D. (2017). A review of web-based chronic disease self-management for older adults. *Gerontechnology*, 16(1), 12–20. <https://doi.org/10.4017/gt.2017.16.1.002.00>
- Poscia A, Stojanovic J, La Milia D I, Duplaga M, Grysztar M, Moscato U, Graziano O, Collamati A, Ricciardi W, & Magnavita N. (2018). Interventions targeting loneliness and social isolation among the older people: An update systematic review. *Experimental Gerontology*, 102, 133–144. <https://doi.org/10.1016/j.exger.2017.11.017>
- Pu L, Moyle W, Jones C, & Todorovic M. (2019). The Effectiveness of Social Robots for Older Adults: A Systematic Review and Meta-Analysis of Randomized Controlled Studies. *Gerontologist*, 59(1), e37–e51. <https://doi.org/10.1093/geront/gny046>
- Queiros A, Santos M, Rocha N P, Cerqueira M, & Ieee. (2017). Technologies for Ageing in Place to Support Community-Dwelling Older Adults. In 2017 12th Iberian Conference on Information Systems and Technologies (*CISTI*), 2017, pp. 1-6, doi: 10.23919/CISTI.2017.7975990.
- Quinn Kelly. (2021). Social media and social wellbeing in later life. *Ageing and Society*, 41(6), 1349–1370. <https://doi.org/10.1017/S0144686X19001570>
- Rainero I, Summers M J, Monter M, Bazzani M, Giannouli E, Aumayr G, Burin D, Provero P, Vercelli A E, & My A H. A. Consortium. (2021). The My Active and Healthy Aging ICT platform prevents quality of life decline in older adults: A randomised controlled study. *Age and Ageing*, 50(4), 1261–1267. <https://doi.org/10.1093/ageing/afaa290>
- Rienzo A, Cubillos C, & Ieee. (2019). Research of Gamification Techniques and their Application in Digital Games for Older Adults. <https://ieeexplore.ieee.org/document/8988019>
- Robbins Ruth E. (2019). The effect of social interaction on isolated older adults. Dissertation Abstracts International: Section B: The Sciences and Engineering, 80(1-B(E)).
<https://repository.arizona.edu/handle/10150/630189>

Robinson H, Macdonald B, Kerse N, & Broadbent E. (2013). The psychosocial effects of a companion robot: A randomized controlled trial. *Journal of the American Medical Directors Association*, 14(9), 661–667. <https://doi.org/10.1016/j.jamda.2013.02.007>

Robinson N L, Cottier T V, & Kavanagh D J. (2019). Psychosocial Health Interventions by Social Robots: Systematic Review of Randomized Controlled Trials. *Journal of Medical Internet Research*, 21(5), e13203. <https://doi.org/10.2196/13203>

Ronzi S, Orton L, Pope D, Valtorta N K, & Bruce N G. (2018). What is the impact on health and wellbeing of interventions that foster respect and social inclusion in community-residing older adults? A systematic review of quantitative and qualitative studies. *Systematic Reviews*, 7(1), 26. <https://doi.org/10.1186/s13643-018-0680-2>

Roth A. (2014). Engaging older cancer patients in psychotherapy and psychoeducation interventions: Care and vintage readers programs. *Psycho-Oncology*, 23, 31.

Schulz Richard, Lustig Amy, Handler Steven, & Martire Lynn M. (2002). Technology-based caregiver intervention research: Current status and future directions. *Gerontechnology*, 2(1), 15–47.

Schwindenhammer Theresa M. (2014). Videoconferencing intervention for depressive symptoms and loneliness in nursing home elders. <https://www.proquest.com/docview/1548716414?pq-origsite=gscholar&fromopenview=true>

Sciamanna C, Ballentine N H, Bopp M, Chinchilli V M, Ciccolo J T, Delauter G, Fisher A, Fox E J, Jan De Beur S, Kearcher K, Kraschnewski J L, Lehman E, McTigue K M, McAuley E, Paranjape A, Rodriguez-Colon S, Rovniak L, Rutt K, Smyth J M, ... Tsay A. (2021). Working to Increase Stability through Exercise (WISE): Screening, recruitment, and baseline characteristics. *Trials*, 22, 1–12. <https://doi.org/10.1186/s13063-022-06018-0>

- Scoglio A A, Reilly E D, Gorman J A, & Drebing C E. (2019). Use of Social Robots in Mental Health and Well-Being Research: Systematic Review. *Journal of Medical Internet Research*, 21(7), e13322. <https://doi.org/10.2196/13322>
- Shah S G. S, Nogueras D, van Woerden H, & Kiparoglou V. (2021). Evaluation of the Effectiveness of Digital Technology Interventions to Reduce Loneliness in Older Adults: Systematic Review and Meta-analysis. *Journal of Medical Internet Research*, 23(6), e24712. <https://doi.org/10.2196/24712>
- Shapira N, Barak A, & Gal I. (2007). Promoting older adults' well-being through Internet training and use. *Aging & Mental Health*, 11(5), 477–484.
- Shapira S, Yeshua-Katz D, Cohn-Schwartz E, Aharonson-Daniel L, Sarid O, & Clarfield A M. (2021). A pilot randomized controlled trial of a group intervention via Zoom to relieve loneliness and depressive symptoms among older persons during the COVID-19 outbreak. *Internet Interventions*, 24, 100368. <https://doi.org/10.1016/j.invent.2021.100368>
- Shapira S, Yeshua-Katz D, Goren G, Aharonson-Daniel L, Clarfield A M, & Sarid O. (2021). Evaluation of a Short-Term Digital Group Intervention to Relieve Mental Distress and Promote Well-Being Among Community-Dwelling Older Individuals During the COVID-19 Outbreak: A Study Protocol. *Frontiers in Public Health*, 9, 577079. <https://doi.org/10.3389/fpubh.2021.577079>
- Shishehgar M, Kerr D, & Blake J. (2018). A systematic review of research into how robotic technology can help older people. *Smart Health*, 7–8, 1–18.
- Shishehgar Majid, Kerr Donald, & Blake Jacqueline. (2019). The effectiveness of various robotic technologies in assisting older adults. *Health Informatics Journal*, 25(3), 892–918. <https://doi.org/10.1177/1460458217729729>
- Smallfield S & Molitor W L. (2018). Occupational Therapy Interventions Supporting Social Participation and Leisure Engagement for Community-Dwelling Older Adults: A

Systematic Review. *American Journal of Occupational Therapy*, 72(4), 7204190020p1-7204190020p8. <https://doi.org/10.5014/ajot.2018.030627>

Smith Lester Jerome. (2011). The effect of training in internet use on perceived loneliness, depression and well-being with the African American elderly. *Dissertation Abstracts International: Section B: The Sciences and Engineering*, 71(12-B), 7354.

Smith Sarah Kate & Astell Arlene. (2017). Independent Living Functions for the Elderly (IN-LIFE): Supporting Communication in Dementia. 242(NA), 16–22.

Smith T L & Toseland R W. (2006). The effectiveness of a telephone support program for caregivers of frail older adults. *Gerontologist*, 46(5), 620–629.

Song J H. (2009). [Effects of a robot pet-assisted program for elderly people with dementia]. *Journal of Korean Academy of Nursing*, 39(4), 562–573.
<https://doi.org/10.4040/jkan.2009.39.4.562>

Song Y & van der Cammen T J. M. (2019). Electronic assistive technology for community-dwelling solo-living older adults: A systematic review. *Maturitas*, 125, 50–56.
<https://doi.org/10.1016/j.maturitas.2019.04.211>

Sosa Giovanni W. (2012). The impact of a video game intervention on the cognitive functioning, self- efficacy, self-esteem, and video game attitudes of older adults. *Dissertation Abstracts International Section A: Humanities and Social Sciences*, 73(5-A), 1885.

Su Z, Meyer K, Li Y, McDonnell D, Joseph N M, Li X, Du Y, Advani S, Cheshmehzangi A, Ahmad J, da Veiga C P, Chung R Y, & Wang J. (2020). Technology-Based Interventions for Nursing Home Residents: Implications for Nursing Home Practice Amid and Beyond the Influence of COVID-19: A Systematic Review Protocol. *Research Square*, 18, 18.
<https://doi.org/10.21203/rs.3.rs-56102/v1>

Tanaka M, Ishii A, Yamano E, Ogikubo H, Okazaki M, Kamimura K, Konishi Y, Emoto S, & Watanabe Y. (2012). Effect of a human-type communication robot on cognitive function in elderly women living alone. *Medical Science Monitor*, 18(9), CR550-7.

- The application of innovative technologies in elderly health care in the community. (2011). *Cyprus Nursing Chronicles*, 12(2), 7–17.
- Thodberg K, Sorensen L U, Christensen J W, Poulsen P H, Houbak B, Damgaard V, Keseler I, Edwards D, & Videbech P B. (2016). Therapeutic effects of dog visits in nursing homes for the elderly. *Psychogeriatrics: The Official Journal of the Japanese Psychogeriatric Society*, 16(5), 289–297. <https://doi.org/10.1111/psyg.12159>
- Tomasino K N, Lattie E G, Ho J, Palac H L, Kaiser S M, & Mohr D C. (2017). Harnessing Peer Support in an Online Intervention for Older Adults with Depression. *American Journal of Geriatric Psychiatry*, 25(10), 1109–1119. <https://doi.org/10.1016/j.jagp.2017.04.015>
- Tsai H H, Cheng C Y, & Shieh W Y. (2020). Effectiveness of laptop-based versus smartphone-based videoconferencing interaction on loneliness, depression and social support in nursing home residents: A secondary data analysis. *Journal of Telemedicine & Telecare*, 1357633X20972004. <https://doi.org/10.1177/1357633X20972004>
- Tsai H H, Cheng C Y, Shieh W Y, & Chang Y C. (2020). Effects of a smartphone-based videoconferencing program for older nursing home residents on depression, loneliness, and quality of life: A quasi-experimental study. *BMC Geriatrics*, 20(1), 27. <https://doi.org/10.1186/s12877-020-1426-2>
- Tsai H H & Tsai Y F. (2011). Changes in depressive symptoms, social support, and loneliness over 1 year after a minimum 3-month videoconference program for older nursing home residents. *Journal of Medical Internet Research*, 13(4), e93. <https://doi.org/10.2196/jmir.1678>
- Tsai H H, Tsai Y F, Wang H H, Chang Y C, & Chu H H. (2010). Videoconference program enhances social support, loneliness, and depressive status of elderly nursing home residents. *Aging & Mental Health*, 14(8), 947–954. <https://doi.org/10.1080/13607863.2010.501057>

- Tyack Charles S. M & Camic Paul M. (2017). Touchscreen interventions and the well-being of people with dementia and caregivers: A systematic review. 29(8), 1261–1280.
<https://doi.org/10.1017/s1041610217000667>
- University of Texas at & Austin. (2020). Telephone Calls for Health for Homebound Older Adults. Clinicaltrials.Gov. <https://clinicaltrials.gov/ct2/show/NCT04595708>
- Uppsala University. (2020). Treatment of Depressive Symptoms in Older Individuals in Isolation During Covid-19. Clinicaltrials.Gov.
<https://clinicaltrials.gov/ct2/show/NCT04508868>
- Van Den Heuvel R, Lexis M, & Daniels R. (2019). CRDL - Interactive technology eliciting engagement in elderly people with dementia. *Technology and Disability*, 31 (Supplement 1), S114.
- Van Houwelingen-Snippe J, Ben Allouch, S, & Van Rompay T J. L. (2021). Virtual Reality Representations of Nature to Improve Well-Being amongst Older Adults: A Rapid Review. *Journal of Technology in Behavioral Science*, 1–22.
<https://doi.org/10.1007/s41347-021-00195-6>
- Vanova M, Irazoki E, Garcia-Casal J A, Martinez-Abad F, Botella C, Shiells K R, & Franco-Martin M A. (2018). The effectiveness of ICT-based neurocognitive and psychosocial rehabilitation programmes in people with mild dementia and mild cognitive impairment using GRADIOR and ehcoBUTLER: study protocol for a randomised controlled trial. *Trials [Electronic Resource]*, 19(1), 100. <https://doi.org/10.1186/s13063-017-2371-z>
- VanRavenstein K, Brotherton S, & Davis B. (2020). Investigating the Feasibility of Using Telemedicine to Deliver a Fall Prevention Program: A Pilot Study. *Journal of Allied Health*, 49(3), 221–227.
- Vazquez F L, Torres A J, Otero P, Blanco V, Lopez L, Garcia-Casal A, & Arrojo M. (2019). Cognitive-behavioral intervention via interactive multimedia online video game for active aging: Study protocol for a randomized controlled trial. *Trials [Electronic Resource]*, 20(1), 692. <https://doi.org/10.1186/s13063-019-3859-5>

- Wasilewski M B, Stinson J N, & Cameron J I. (2017). Web-based health interventions for family caregivers of elderly individuals: A Scoping Review. *International Journal of Medical Informatics*, 103, 109–138. <https://doi.org/10.1016/j.ijmedinf.2017.04.009>
- White H, McConnell E, Clipp E, Branch L G, Sloane R, Pieper C, & Box T L. (2002). A randomized controlled trial of the psychosocial impact of providing internet training and access to older adults. *Aging & Mental Health*, 6(3), 213–221. <https://doi.org/10.1080/13607860220142422>
- White Heidi K, McConnell Eleanor S, Clipp Elizabeth C, Bynum Louise, Teague Carmen, Navas Luis, Craven Sara, & Halbrecht Herbert. (1999). Surfing the Net in Later Life: A Review of the Literature and Pilot Study of Computer Use and Quality of Life. 18(3), 358–378. <https://doi.org/10.1177/073346489901800306>
- Woodward A T, Freddolino P P, Blaschke-Thompson C M, Wishart D J, Bakk L, Kobayashi R, & Tupper C. (2011). Technology and Aging Project: Training Outcomes and Efficacy from a Randomized Field Trial. *Ageing International*, 36(1), 46–65. <https://doi.org/10.1007/s12126-010-9074-z>
- Zaccaria D, Guaita A, Vaccaro R, Casanova G, Abbondanza S, Pettinato L, Cerati G, Rolandi E, & Sala E. (2020). Assessing the impact of Social Networking Site use on older people's loneliness and social isolation. A randomized controlled trial: The Aging in a Networked Society-Social Experiment Study (ANS-SE). *Contemporary Clinical Trials Communications*, 19, 100615. <https://doi.org/10.1016/j.conctc.2020.100615>
- Zhao Xia, Wang LiNa, Ge ChenXi, Zhen XueTing, Chen Zheng, Wang Jie, & Zhou YuanYuan. (2020). Smartphone application training program improves smartphone usage competency and quality of life among the elderly in an elder university in China: A randomized controlled trial. *International Journal of Medical Informatics*, 133. <https://doi.org/10.1016/j.ijmedinf.2019.104010>

Excluded studies

- Bolle, S., van Weert, J. C., Daams, J. G., Loos, E. F., De Haes, H. C., & Smets, E. M. (2015). Online health information tool effectiveness for older patients: a systematic review of the literature. *Journal of health communication*, 20(9), 1067-1083.
- Burkow, T. M., Vognild, L. K., Johnsen, E., Risberg, M. J., Bratvold, A., Breivik, E., ... & Hjalmsarsen, A. (2015). Comprehensive pulmonary rehabilitation in home-based online groups: a mixed method pilot study in COPD. *BMC research notes*, 8(1), 1-11.
- Clarkson, P., Hughes, J., Roe, B., Giebel, C. M., Jolley, D., Poland, F., ... & Challis, D. (2018). Systematic review: Effective home support in dementia care, components and impacts—Stage 2, effectiveness of home support interventions. *Journal of Advanced Nursing*, 74(3), 507-527.
- Cooper, K., Kirkpatrick, P., & Wilcock, S. (2014). The effectiveness of peer support interventions for community-dwelling adults with chronic non-cancer pain: A systematic review. *JBI Evidence Synthesis*, 12(5), 319-348.
- Dickens, A. P., Richards, S. H., Hawton, A., Taylor, R. S., Greaves, C. J., Green, C., ... & Campbell, J. L. (2011). An evaluation of the effectiveness of a community mentoring service for socially isolated older people: a controlled trial. *BMC Public Health*, 11(1), 1-14.
- Dam, A. E., de Vugt, M. E., van Boxtel, M. P., & Verhey, F. R. (2017). Effectiveness of an online social support intervention for caregivers of people with dementia: the study protocol of a randomised controlled trial. *Trials*, 18(1), 1-10.
- Erfani, S. S., & Abedin, B. (2018). Impacts of the use of social network sites on users' psychological well-being: A systematic review. *Journal of the Association for Information Science and Technology*, 69(7), 900-912.
- Fan, Q. (2016). Utilizing ICT to prevent loneliness and social isolation of the elderly: a literature review. *Cuadernos de Trabajo Social*, 29(2).

- Forsman, A. K., & Nordmyr, J. (2017). Psychosocial links between Internet use and mental health in later life: a systematic review of quantitative and qualitative evidence. *Journal of Applied Gerontology*, 36(12), 1471-1518.
- Giné-Garriga, M., Coll-Planas, L., Guerra, M., Domingo, À., Roqué, M., Caserotti, P., ... & Salvà, A. (2017). The SITLESS project: exercise referral schemes enhanced by self-management strategies to battle sedentary behaviour in older adults: study protocol for a randomised controlled trial. *Trials*, 18(1), 1-14.
- Gorenko, J. A., Moran, C., Flynn, M., Dobson, K., & Konnert, C. (2021). Social isolation and psychological distress among older adults related to COVID-19: a narrative review of remotely-delivered interventions and recommendations. *Journal of Applied Gerontology*, 40(1), 3-13.
- Jones, C. A., Siever, J., Knuff, K., Van Bergen, C., Mick, P., Little, J., ... & Miller, H. (2019). Walk, Talk and Listen: a pilot randomised controlled trial targeting functional fitness and loneliness in older adults with hearing loss. *BMJ open*, 9(4), e026169.
- Lara, J., O'Brien, N., Godfrey, A., Heaven, B., Evans, E. H., Lloyd, S., ... & Mathers, J. C. (2016). Pilot randomised controlled trial of a web-based intervention to promote healthy eating, physical activity and meaningful social connections compared with usual care control in people of retirement age recruited from workplaces. *PloS one*, 11(7), e0159703.
- Nicholson, N. R. (2012). A review of social isolation: an important but underassessed condition in older adults. *The journal of primary prevention*, 33(2), 137-152.
- Nijman, S. A., Veling, W., Greaves-Lord, K., Vermeer, R. R., Vos, M., Zandee, C. E., ... & Pijnenborg, G. H. (2019). Dynamic Interactive Social Cognition Training in Virtual Reality (DiSCoVR) for social cognition and social functioning in people with a psychotic disorder: study protocol for a multicenter randomized controlled trial. *BMC psychiatry*, 19(1), 1-11.

- Perkins, E. A., & LaMartin, K. M. (2012). The Internet as social support for older carers of adults with intellectual disabilities. *Journal of Policy and Practice in Intellectual Disabilities*, 9(1), 53-62.
- Preston, C., & Moore, S. (2019). Ringing the changes: the role of telephone communication in a helpline and befriending service targeting loneliness in older people. *Ageing & Society*, 39(7), 1528-1551.
- Rebollar, A. M., & Francisco, W. C. (2015). Correlación entre actividades de interacción social registradas con nuevas tecnologías y el grado de aislamiento social en los adultos mayores. *Revista mexicana de ingeniería biomédica*, 36(3), 181-190.
- Selak, Š., Bacaicoa, O. A., & Gabrovec, B. (2019). Can we manage frailty at individual level by the use of information and communication technologies: A narrative literature review. *Slovenian Medical Journal*, 88(5-6), 249-262.
- Sumner, J., Chong, L. S., Bundele, A., & Wei Lim, Y. (2021). Co-designing technology for aging in place: A systematic review. *The Gerontologist*, 61(7), e395-e409.
- Toh, H. M., Ghazali, S. E., & Subramaniam, P. (2016). The acceptability and usefulness of cognitive stimulation therapy for older adults with dementia: A narrative review. *International journal of Alzheimer's disease*, 2016.
- Winterton, R., & Warburton, J. (2011). Models of care for socially isolated older rural carers: barriers and implications. *Rural and Remote Health*, 11(3), 203-215.
- Zeppegno, P., Gattoni, E., Mastrangelo, M., Gramaglia, C., & Sarchiapone, M. (2019). Psychosocial suicide prevention interventions in the elderly: a mini-review of the literature. *Frontiers in psychology*, 2713.

Other references

Additional references

- Abdi, S., Spann, A., Borilovic, J., de Witte, L., & Hawley, M. (2019). Understanding the care and support needs of older people: A scoping review and categorisation using the WHO

- international classification of functioning, disability and health framework (ICF). *BMC Geriatrics*, 19(1), 1–15.
- Bedney, B. J., Goldberg, R. B., & Josephson, K. (2010). Aging in place in naturally occurring retirement communities: Transforming aging through supportive service programs. *Journal of Housing for the Elderly*, 24(3–4), 304–321.
- Berkman, L. F., Glass, T., Brissette, I., & Seeman, T. E. (2000). From social integration to health: Durkheim in the new millennium. *Social Science & Medicine*, 51(6), 843–857.
- Bethell, J., Babineau, J., Iaboni, A., Green, R., Cuaresma-Canlas, R., Karunanathan, R., Schon, B., Schon, D., & McGilton, K. S. (2019). Social integration and loneliness among long-term care home residents: Protocol for a scoping review. *BMJ Open*, 9(12), e033240.
- Boulton, E., Kneale, D., Stansfield, C., Heron, P., Sutcliffe, K., Hayanga, B., Hall, A., Bower, P., Casey, D., & Craig, D. (2020). Rapid systematic review of systematic reviews: What befriending, social support and low intensity psychosocial interventions, delivered remotely, are effective in reducing social isolation and loneliness among older adults? How do they work?[version 1; peer review: Awaiting peer review]. *F1000Research*, 9.
- Brooke, J., & Jackson, D. (2020). Older people and COVID-19: Isolation, risk and ageism. *Journal of Clinical Nursing*, 29(13–14), 2044–2046.
- Budd, J., Miller, B. S., Manning, E. M., Lampos, V., Zhuang, M., Edelstein, M., Rees, G., Emery, V. C., Stevens, M. M., & Keegan, N. (2020). Digital technologies in the public-health response to COVID-19. *Nature Medicine*, 26(8), 1183–1192.
- Cattan, M., White, M., Bond, J., & Learmouth, A. (2005). Preventing social isolation and loneliness among older people: A systematic review of health promotion interventions. *Ageing & Society*, 25(1), 41–67.
- Chen, Y.-R. R., & Schulz, P. J. (2016). The effect of information communication technology interventions on reducing social isolation in the elderly: A systematic review. *Journal of Medical Internet Research*, 18(1), e18.
- Chipps, J., Jarvis, M. A., & Ramlall, S. (2017). The effectiveness of e-Interventions on reducing social isolation in older persons: A systematic review of systematic reviews. *Journal of*

Telemedicine and Telecare, 23(10), 817–827.

<https://doi.org/10.1177/13576333X17733773>

- Cohen-Mansfield, J., & Perach, R. (2015). Interventions for alleviating loneliness among older persons: A critical review. *American Journal of Health Promotion*, 29(3), e109–e125.
- DeGood, K. (2011). *Aging in place, stuck without options: Fixing the mobility crisis threatening the baby boom generation*.
- Dickens, A. P., Richards, S. H., Greaves, C. J., & Campbell, J. L. (2011). Interventions targeting social isolation in older people: A systematic review. *BMC Public Health*, 11(1), 1–22.
- Digital Solution Foundry and EPPI_Centre. (2020). *Eppi-Mapper* (1.2.5) [Computer software].
- Donovan, N. J., & Blazer, D. (2020). Social isolation and loneliness in older adults: Review and commentary of a National Academies Report. *The American Journal of Geriatric Psychiatry*, 28(12), 1233–1244.
- Fakoya, O. A., McCorry, N. K., & Donnelly, M. (2020). Loneliness and social isolation interventions for older adults: A scoping review of reviews. *BMC Public Health*, 20(1), 1–14.
- Falk. (2021). *The state of virtual care in Canada as of wave three of the COVID-19 pandemic: An Early Diagnostic and Policy Recommendations*. Health Canada.
https://www.canada.ca/content/dam/hc-sc/documents/corporate/transparency_229055456/health-agreements/bilateral-agreement-pan-canadian-virtual-care-priorities-covid-19/template-wf-report-eng.pdf
- Findlay, R. A. (2003). Interventions to reduce social isolation amongst older people: Where is the evidence? *Ageing & Society*, 23(5), 647–658.
- Gardiner, C., Geldenhuys, G., & Gott, M. (2018). Interventions to reduce social isolation and loneliness among older people: An integrative review. *Health & Social Care in the Community*, 26(2), 147–157.

- Hagan, R., Manktelow, R., Taylor, B. J., & Mallett, J. (2014). Reducing loneliness amongst older people: A systematic search and narrative review. *Aging & Mental Health, 18*(6), 683–693.
- Henderson, E. J., & Caplan, G. A. (2008). Home sweet home? Community care for older people in Australia. *Journal of the American Medical Directors Association, 9*(2), 88–94.
- Holt-Lunstad, J., Smith, T. B., Baker, M., Harris, T., & Stephenson, D. (2015). Loneliness and social isolation as risk factors for mortality: A meta-analytic review. *Perspectives on Psychological Science, 10*(2), 227–237.
- Ibarra, F., Baez, M., Cernuzzi, L., & Casati, F. (2020). A systematic review on technology-supported interventions to improve old-age social wellbeing: Loneliness, social isolation, and connectedness. *Journal of Healthcare Engineering, 2020*.
- Jopling, K. (2020). Promising approaches revisited: Effective action on loneliness in later life. *Campaign to End Loneliness*.
- Kelly, M. E., Duff, H., Kelly, S., Power, J. E. M., Brennan, S., Lawlor, B. A., & Loughrey, D. G. (2017). The impact of social activities, social networks, social support and social relationships on the cognitive functioning of healthy older adults: A systematic review. *Systematic Reviews, 6*(1), 1–18.
- Khosravi, P., Rezvani, A., & Wiewiora, A. (2016). The impact of technology on older adults' social isolation. *Computers in Human Behavior, 63*, 594–603.
<https://doi.org/10.1016/j.chb.2016.05.092>
- Leigh-Hunt, N., Bagguley, D., Bash, K., Turner, V., Turnbull, S., Valtorta, N., & Caan, W. (2017). An overview of systematic reviews on the public health consequences of social isolation and loneliness. *Public Health, 152*, 157–171.
- Mann, F., Bone, J. K., Lloyd-Evans, B., Frerichs, J., Pinfold, V., Ma, R., Wang, J., & Johnson, S. (2017). A life less lonely: The state of the art in interventions to reduce loneliness in people with mental health problems. *Social Psychiatry and Psychiatric Epidemiology, 52*(6), 627–638.

- Masi, C. M., Chen, H.-Y., Hawkley, L. C., & Cacioppo, J. T. (2011). A meta-analysis of interventions to reduce loneliness. *Personality and Social Psychology Review, 15*(3), 219–266.
- Menec, V. H., Newall, N. E., Mackenzie, C. S., Shooshtari, S., & Nowicki, S. (2020). Examining social isolation and loneliness in combination in relation to social support and psychological distress using Canadian Longitudinal Study of Aging (CLSA) data. *PLOS ONE, 15*(3), e0230673.
- Moher, D., Shamseer, L., Clarke, M., Ghersi, D., Liberati, A., Petticrew, M., Shekelle, P., & Stewart, L. A. (2015). Preferred reporting items for systematic review and meta-analysis protocols (PRISMA-P) 2015 statement. *Systematic Reviews, 4*(1), 1–9.
- Newall, N. E., & Menec, V. H. (2019). Loneliness and social isolation of older adults: Why it is important to examine these social aspects together. *Journal of Social and Personal Relationships, 36*(3), 925–939.
- Poscia, A., Stojanovic, J., La Milia, D. I., Duplaga, M., Grysztar, M., Moscato, U., Onder, G., Collamati, A., Ricciardi, W., & Magnavita, N. (2018). Interventions targeting loneliness and social isolation among the older people: An update systematic review. *Experimental Gerontology, 102*, 133–144.
- Prohaska, T., Burholt, V., Burns, A., Golden, J., Hawkley, L., Lawlor, B., Leavey, G., Lubben, J., O’Sullivan, R., & Perissinotto, C. (2020). Consensus statement: Loneliness in older adults, the 21st century social determinant of health? *BMJ Open, 10*(8), e034967.
- Shah, S. G. S., Nogueras, D., Van Woerden, H. C., & Kiparoglou, V. (2020). The COVID-19 pandemic: A pandemic of lockdown loneliness and the role of digital technology. *Journal of Medical Internet Research, 22*(11), e22287.
- Shah, S. G. S., Nogueras, D., van Woerden, H. C., & Kiparoglou, V. (2021). Evaluation of the effectiveness of digital technology interventions to reduce loneliness in older adults: Systematic review and meta-analysis. *Journal of Medical Internet Research, 23*(6), e24712.

- Shea, B. J., Reeves, B. C., Wells, G., Thuku, M., Hamel, C., Moran, J., Moher, D., Tugwell, P., Welch, V., & Kristjansson, E. (2017). AMSTAR 2: A critical appraisal tool for systematic reviews that include randomised or non-randomised studies of healthcare interventions, or both. *Bmj*, *358*.
- Snilstveit, B., Vojtkova, M., Bhavsar, A., & Gaarder, M. (2013). *Evidence gap maps—A tool for promoting evidence-informed policy and prioritizing future research*. The world bank.
- Snilstveit, B., Vojtkova, M., Bhavsar, A., Stevenson, J., & Gaarder, M. (2016). Evidence & Gap Maps: A tool for promoting evidence informed policy and strategic research agendas. *Journal of Clinical Epidemiology*, *79*, 120–129.
- ten Bruggencate, T., Luijkx, K. G., & Sturm, J. (2019). How to fulfil social needs of older people: Exploring design opportunities for technological interventions. *Gerontechnology*, *18*(3), 156–167.
- Thangavel, G., Memedi, M., & Hedström, K. (2022). Customized Information and Communication Technology for Reducing Social Isolation and Loneliness Among Older Adults: Scoping Review. *JMIR Mental Health*, *9*(3), e34221.
- Thomas, J., Graziosi, S., Brunton, J., Ghouze, Z., O’Driscoll, P., & Bond, M. (2020). EPPI-Reviewer: Advanced software for systematic reviews, maps and evidence synthesis. *Eppi-Centre Software*. London: UCL Social Research Institute.
- Tomaka, J., Thompson, S., & Palacios, R. (2006). The relation of social isolation, loneliness, and social support to disease outcomes among the elderly. *Journal of Aging and Health*, *18*(3), 359–384.
- UCLG. (2020). *Live Learning Experience: Beyond the immediate response to the outbreak of COVID-19. Digital Technologies and the COVID-19 pandemic. Briefing & Learning Note 15.04, 2020*. United Cities and Local Governments (UCLG).
https://www.uclg.org/sites/default/files/eng_briefing_technology_final_x.pdf
- Victor, C., Mansfield, L., Kay, T., Daykin, N., Lane, J., Grigsby Duffy, L., Tomlinson, A., & Meads, C. (2018). An overview of reviews: The effectiveness of interventions to address loneliness at all stages of the life-course. *What Works Centre for Wellbeing*.

- Victor, C. R., Scambler, S., Bowling, A. N. N., & Bond, J. (2005). The prevalence of and risk factors for, loneliness in later life: A survey of older people in Great Britain. *Ageing and Society*, 25(6), 357–375.
- Waddington, H., Snilstveit, B., Hombrados, J., Vojtkova, M., Phillips, D., Davies, P., & White, H. (2014). Farmer field schools for improving farming practices and farmer outcomes: A systematic review. *Campbell Systematic Reviews*, 10(1), i–335.
- Watts, G. (2020). COVID-19 and the digital divide in the UK. *The Lancet Digital Health*, 2(8), e395–e396.
- Welch, V., Ghogomu, E.T., Barbeau, V.I., Boulton, E., Boutin, S., Haitas, N., Kneale, D., Salzwedel, D. M, Simard, R., Herbert, P., & Mikton, C. (2022). PROTOCOL: Digital interventions to reduce social isolation and loneliness in older adults: An evidence and gap map. *Campbell Systematic Reviews*, 18(3), e1260.
- White, H., Albers, B., Gaarder, M., Kornør, H., Littell, J., Marshall, Z., Matthew, C., Pigott, T., Snilstveit, B., & Waddington, H. (2020). Guidance for producing a Campbell evidence and gap map. *Campbell Systematic Reviews*, 16(4), e1125.
- Williams, C. Y. K., Townson, A. T., Kapur, M., Ferreira, A. F., Nunn, R., Galante, J., Phillips, V., Gentry, S., & Usher-Smith, J. A. (2021). Interventions to reduce social isolation and loneliness during COVID-19 physical distancing measures: A rapid systematic review. *PLOS ONE*, 16(2), e0247139. <https://doi.org/10.1371/journal.pone.0247139>
- Windle, K., Francis, J., & Coomber, C. (2011). *Preventing loneliness and social isolation: Interventions and outcomes. SCIE Research Briefing 39*. London: Social Care Institute for Excellence, 1–16.
- World Bank. (2021). *World Bank country and lending groups. Country classification*. World Bank Data Help Desk.
- World Health Organization. (2018). *Classification of digital health interventions v1. 0: A shared language to describe the uses of digital technology for health*. World Health Organization.
- World Health Organization. (2019). *World health statistics 2019: Monitoring health for the SDGs, sustainable development goals*. World Health Organization.

World Health Organization. (2020). *Decade of healthy ageing: Baseline report*.

World Health Organization. (2021). *Social isolation and loneliness among older people: Advocacy brief*.

Zanella, A., Mason, F., Pluchino, P., Cisotto, G., Orso, V., & Gamberini, L. (2020). Internet of things for elderly and fragile people. *ArXiv Preprint ArXiv:2006.05709*.

Zavaleta, D., Samuel, K., & Mills, C. (2014). *OPHI Working Paper No. 67. Social isolation: A conceptual and measurement proposal*. Oxford Poverty & Human Development Initiative, University of Oxford: Oxford, UK.

Sources of support

Internal sources

- None

External sources

- World Health Organization funding - Purchase Order Number: 202666968

Appendices

1. Glossary of key concepts

Key concepts related to intervention categories

Interventions to improve social skills: Interventions that focus on training in or education on one's social skills, such as conversational ability. They aim to enable individuals to form and maintain meaningful relationships (Mann et al., 2017; Masi et al., 2011).

Social skills are any skills facilitating interaction and communication with others. They are socially acceptable learned behaviours that enable a person to interact with others in ways that elicit positive responses and assist in avoiding negative responses (Little et al., 2017).

Interventions to enhance social support: interventions where people are offered support (regular contacts, care, or companionship) and guidance in finding and attending new

activities or groups. They aim to help individuals make and maintain social connections (Mann et al., 2017; Masi et al., 2011).

Social support is the actual or perceived availability of resources (e.g., tangible, informational, emotional help) from others, typically one's social network (Berkman et al., 2000; Bethell et al., 2019; Kelly et al., 2017).

Interventions to enhance social interactions: interventions that focus on improving the quality of relationships and increase opportunities for social interactions. They aim to promote engagement (Masi et al., 2011).

Interventions for social cognitive training: interventions that focus on changing negative thinking and feelings about social relationships. They aim to change behaviours, reduce maladaptive cognitions and increase social connections (Mann et al., 2017; Masi et al., 2011).

Key concepts related to outcomes

Social connection: encompasses the structural, functional, and quality aspects of how individuals connect to each other (Donovan & Blazer, 2020; Zavaleta et al., 2014).

Social network: denotes the number and type of social relationships with people (individuals or groups) (Berkman et al., 2000; Bethell et al., 2019; Kelly et al., 2017; Mann et al., 2017).

Social support: the actual or perceived availability of resources (e.g., tangible, informational, emotional help) from others, typically one's social network (Berkman et al., 2000; Bethell et al., 2019; Kelly et al., 2017; Mann et al., 2017) to help them adapt to change and cope with stress (Elder & Retrum, 2012).

Social engagement: reflects participation in meaningful activities with others (Berkman et al., 2000; Bethell et al., 2019; Elder & Retrum, 2012).

Social cohesion: is the neighborhood-based resource of mutual community trust and solidarity (Elder & Retrum, 2012).

Social capital: refers to a series of resources that individuals earn as a result of their membership in social networks, and the features of those networks that facilitate individual or collective actions such as interpersonal trust, reciprocity and mutual aid (Bethell et al., 2019; Elder & Retrum, 2012; Mann et al., 2017; Zavaleta et al., 2014).

Digital divide: refers to disparities in access to technological interventions (e.g., smartphones, computers and the internet) which may be due to lack of affordability or access to technology, broadband or Wi-Fi, data poverty or geographic location, lack of digital skills and the confidence to access online services and support (Budd et al., 2020; UCLG, 2020; Watts, 2020).

Other concepts

Evidence architecture: The system by which evidence to inform decision-making is produced and made available to decision-makers. The system spans data collection and data collection infrastructure (such as initiatives to standardize outcome indicators), the analysis and reporting of those data in studies, and summaries of such studies in systematic reviews, through to products with varying degrees of curation such as evidence platforms, guidelines and checklists. (White, 2019, 2020, 2021).

2. Search strategies

Database: Ovid MEDLINE(R) ALL <1946 to May 14, 2021>

Search Date: 16 May 2021

Yield: 4685

-
- 1 exp aged/ or middle aged/ (5245060)
 - 2 (ageing or aging or care home resident* or community-dwelling or elder* or frail* or geriatric* or "late life" or "later life" or "mature adult*" or "middle age*" or "mid* life" or midlife or nonagenarian* or nursing home resident* or octogenarian* or old age* or "oldest old" or pensioner* or postmenopaus* or post-menopaus* or retired or retiree* or senior citizen* or seniors or (older adj3 adult*) or old age* or (older adj3 client*) or (older adj3 communit*) or (older adj3 female*) or (older adj3 individual*) or (older adj3 inpatient*) or (older adj3 in-patient*) or (older adj3 male*) or (older adj3 men) or (older adj3 outpatient*) or (older adj3 out-patient*) or (older adj3 patient*) or (older adj3 people) or (older adj3 person*) or (older adj3 population*) or (older adj3 women)).tw,kf. /freq=2 (422994)
 - 3 or/1-2 (5385267)
 - 4 *communication/ or *loneliness/ or *social isolation/ or social support/ (118907)

- 5 (befriend* or connectedness or (connect* adj2 other*) or (connect* adj2 people) or friendship* or interpersonal or loneliness or lonely or make friend* or making friend* or psychosocial or wellbeing or well-being).tw,kf. (256454)
- 6 ((enhanc* or improv* or increas* or maintain*) adj5 (communicat* or connect* or interaction*)).tw,kf. (90243)
- 7 ((social* adj2 contact*) or (social* adj2 connect*) or (social* adj2 distanc*) or (social* adj2 exclu*) or (social* adj2 inclu*) or (social* adj2 interact*) or (social* adj2 isolat*) or (social* adj2 participat*) or (social* adj2 relation*)).tw,kf. (75647)
- 8 or/4-7 (487286)
- 9 technology.hw. (96226)
- 10 *computers/ or *computer user training/ or *mobile applications/ or *social media/ or *user-computer interface/ or *wireless technology/ (57698)
- 11 (app or apps or audio* or chatbot* or chat forum* or chat interface* or chatroom* or chat room* or chat site* or chat software or computer* or digital* or DVD* or email* or e-mail* or electronic or facebook* or facetime or helpline* or help-line* or instant messag* or live chat* or messaging or mms or mobile app* or online* or phone* or phoning or robot* or short messag* service* or skype* or smartphone* or sms or social network* or social media or support line* or technolog* or telephon* or texting or text messag* or tweets or twitter or video conferenc* or videoconferenc* or video record* or videorecord* or virtual* or web app* or web-based or webcam* or webcast* or web portal* or website* or web site* or "web 2.0" or "web 3.0" or whatsapp or wireless or world wide web).ti,kf. (469878)
- 12 (app or apps or audio* or chatbot* or chat forum* or chat interface* or chatroom* or chat room* or chat site* or chat software or computer* or digital* or DVD* or email* or e-mail* or electronic or facebook* or facetime or helpline* or help-line* or instant messag* or live chat* or messaging or mms or mobile app* or online* or phone* or phoning or robot* or short messag* service* or skype* or smartphone* or sms or social network* or social media or support line* or technolog* or telephon* or texting or text messag* or tweets or twitter or video conferenc* or videoconferenc* or video record* or videorecord* or virtual* or web app* or web-based or webcam* or webcast* or web portal* or website* or web site* or "web 2.0" or "web 3.0" or whatsapp or wireless or world wide web).ab. /freq=2 (480103)
- 13 (e-counsel* or ehealth* or e-health or e-intervention* or esupport* or e-support* or etherap* or e-therap* or mhealth* or m-health or mobile health* or online CBT or

- remote care or remote consult* or remote* deliver* or remote health* or self-care or self-guided or self-help or telecare or tele-care or teleconsult* or tele-consult* or telehealth* or tele-health* or telegroup* or tele-group* or telemed* or tele-med* or teletherap* or tele-therap* or virtual realit*).ti,kf. (46974)
- 14 (e-counsel* or ehealth* or e-health or e-intervention* or esupport* or e-support* or etherap* or e-therap* or mhealth* or m-health or mobile health* or remote care or remote consult* or remote* deliver* or remote health* or self-care or self-guided or self-help or telecare or tele-care or teleconsult* or tele-consult* or telehealth* or tele-health* or telegroup* or tele-group* or telemed* or tele-med* or teletherap* or tele-therap* or virtual realit*).ab. /freq=2 (26582)
- 15 or/9-14 (859494)
- 16 (benefit* or change or changes or contribut* or decreas* or develop* or effect or effects or effectiveness or enhance* or evaluat* or experience* or experiment* or impact* or implement* or increas* or intervention* or method* or outcome* or pilot* or program* or provid* or reduc* or study or support* or system* or target* or technolog* or training or trial or "use of ").ti,kf. (9457713)
- 17 systematic review.mp,pt. (214838)
- 18 meta analysis.mp,pt. (213822)
- 19 (cochrane or embase or medline or pubmed).ab. (248403)
- 20 randomized controlled trial.pt. (529685)
- 21 controlled clinical trial.pt. (94148)
- 22 pragmatic clinical trial.pt. (1735)
- 23 randomi*.tw,kf. (688878)
- 24 placebo.ab. (218033)
- 25 clinical trials as topic/ (195808)
- 26 (randomly adj2 (allocated or assigned)).ab. (149132)
- 27 trial.ti. (239911)
- 28 (group or groups).ab. /freq=2 (1953207)
- 29 ((quasi experiment* or quasiexperiment* or quasi randomi* or quasirandomi*) adj2 (design* or method* or study or trial)).ab,kf. (11681)
- 30 ((before adj5 after) or (controlled adj3 study) or (controlled adj3 trial) or control group* or effect? or evaluat* or experience* or impact? or intervention* or (pre adj5 post) or ((pretest or pre-test) and (posttest or post test))).ab,kf. /freq=2 (4635358)
- 31 controlled before-after studies/ (612)

- 32 (controlled before* adj2 after).ab,kf. (1116)
33 interrupted time series analysis/ (1215)
34 (time series adj5 (analys* or design* or interrupted or ITS or studies or study or
trial)).ab,kf. (13874)
35 or/17-34 (6471051)
36 animals/ not (humans/ and animals/) (4791745)
37 35 not 36 (5374053)
38 3 and 8 and 15 and 16 and 37 (4685)
-

Database: Embase Classic+Embase <1947 to 2021 May 14>

Search Date: 16 May 2021

Yield: 4003

- 1 exp aged/ or middle aged/ (4423286)
2 (ageing or aging or care home resident* or community-dwelling or elder* or frail* or
geriatric* or "late life" or "later life" or "mature adult*" or "middle age*" or "mid* life" or
midlife or nonagenarian* or nursing home resident* or octogenarian* or old age* or
"oldest old" or pensioner* or postmenopaus* or post-menopaus* or retired or retiree* or
senior citizen* or seniors or (older adj3 adult*) or old age* or (older adj3 client*) or
(older adj3 communit*) or (older adj3 female*) or (older adj3 individual*) or (older adj3
inpatient*) or (older adj3 in-patient*) or (older adj3 male*) or (older adj3 men) or (older
adj3 outpatient*) or (older adj3 out-patient*) or (older adj3 patient*) or (older adj3
people) or (older adj3 person*) or (older adj3 population*) or (older adj3 women)).tw.
/freq=2 (568588)
3 or/1-2 (4621214)
4 *communication/ or *loneliness/ or *social isolation/ or social support/ (146374)
5 (befriend* or connectedness or (connect* adj2 other*) or (connect* adj2 people) or
friendship* or interpersonal or loneliness or lonely or make friend* or making friend* or
psychosocial or wellbeing or well-being).tw. (328973)
6 ((enhanc* or improv* or increas* or maintain*) adj5 (communicat* or connect* or
interaction*)).tw. (116475)

- 7 ((social* adj2 contact*) or (social* adj2 connect*) or (social* adj2 distanc*) or (social* adj2 exclu*) or (social* adj2 inclu*) or (social* adj2 interact*) or (social* adj2 isolat*) or (social* adj2 participat*) or (social* adj2 relation*)).tw. (93277)
- 8 or/4-7 (619385)
- 9 technology.hw. (198580)
- 10 exp *computers/ or *human computer interaction/ or exp *mobile application/ or *social media/ or *computer interface/ or *wireless communication/ (92346)
- 11 (app or apps or audio* or chatbot* or chat forum* or chat interface* or chatroom* or chat room* or chat site* or chat software or computer* or digital* or DVD* or email* or e-mail* or electronic or facebook* or facetime or helpline* or help-line* or instant messag* or live chat* or messaging or mms or mobile app* or online* or phone* or phoning or robot* or short messag* service* or skype* or smartphone* or sms or social network* or social media or support line* or technolog* or telephon* or texting or text messag* or tweets or twitter or video conferenc* or videoconferenc* or video record* or videorecord* or virtual* or web app* or web-based or webcam* or webcast* or web portal* or website* or web site* or "web 2.0" or "web 3.0" or whatsapp or wireless or world wide web).ti. (499318)
- 12 (app or apps or audio* or chatbot* or chat forum* or chat interface* or chatroom* or chat room* or chat site* or chat software or computer* or digital* or DVD* or email* or e-mail* or electronic or facebook* or facetime or helpline* or help-line* or instant messag* or live chat* or messaging or mms or mobile app* or online* or phone* or phoning or robot* or short messag* service* or skype* or smartphone* or sms or social network* or social media or support line* or technolog* or telephon* or texting or text messag* or tweets or twitter or video conferenc* or videoconferenc* or video record* or videorecord* or virtual* or web app* or web-based or webcam* or webcast* or web portal* or website* or web site* or "web 2.0" or "web 3.0" or whatsapp or wireless or world wide web).ab. /freq=2 (644410)
- 13 (e-counsel* or ehealth* or e-health or e-intervention* or esupport* or e-support* or etherap* or e-therap* or mhealth* or m-health or mobile health* or online CBT or remote care or remote consult* or remote* deliver* or remote health* or self-care or self-guided or self-help or telecare or tele-care or teleconsult* or tele-consult* or telehealth* or tele-health* or telegroup* or tele-group* or telemed* or tele-med* or teletherap* or tele-therap* or virtual realit*).ti. (39628)

- 14 (e-counsel* or ehealth* or e-health or e-intervention* or esupport* or e-support* or
etherap* or e-therap* or mhealth* or m-health or mobile health* or online CBT or
remote care or remote consult* or remote* deliver* or remote health* or self-care or self-
guided or self-help or telecare or tele-care or teleconsult* or tele-consult* or telehealth*
or tele-health* or telegroup* or tele-group* or telemed* or tele-med* or teletherap* or
tele-therap* or virtual realit*).ab. /freq=2 (34397)
- 15 or/9-14 (1089295)
- 16 (benefit* or change or changes or contribut* or decreas* or develop* or effect or effects or
effectiveness or enhance* or evaluat* or experience* or experiment* or impact* or
implement* or increas* or intervention* or method* or outcome* or pilot* or program*
or provid* or reduc* or study or support* or system* or target* or technolog* or training
or trial or "use of").ti. (11478419)
- 17 systematic review.tw. (228178)
- 18 meta analysis*.tw. (222815)
- 19 (cochrane or embase or medline or pubmed).ab. (314793)
- 20 randomized controlled trial/ (658114)
- 21 crossover procedure/ (67193)
- 22 double-blind procedure/ (186270)
- 23 randomi*.tw. (987166)
- 24 (crossover* or cross-over*).tw. (113780)
- 25 placebo.ab. (319158)
- 26 ((doubl* or singl*) adj blind*).tw. (250056)
- 27 (randomly adj2 (allocated or assigned)).ab. (188107)
- 28 (group or groups).ab. /freq=2 (2908853)
- 29 ((quasi experiment* or quasiexperiment* or quasi randomi* or quasirandomi*) adj2
(design* or method* or study or trial)).ab. (14180)
- 30 ((before adj5 after) or (controlled adj3 study) or (controlled adj3 trial) or control group* or
effect? or evaluat* or experience* or impact? or intervention* or (pre adj5 post) or
((pretest or pre-test) and (posttest or post test))).ab. /freq=2 (6397976)
- 31 (controlled before* adj2 after).ab. (1257)
- 32 time series analysis/ (29066)
- 33 (time series adj5 (analys* or design* or interrupted or ITS or studies or study or trial)).ab.
(15610)
- 34 or/17-33 (8761701)

- 35 (exp animal/ or animal.hw. or nonhuman/) not (exp human/ or human cell/ or (human or humans).ti.) (7392778)
- 36 34 not 35 (7091401)
- 37 3 and 8 and 15 and 16 and 36 (4003)
-

Database: APA PsycInfo <1806 to May Week 2 2021> via Ovid

Search Date: 16 May 2021

Yield: 1214

- 1 aged/ or exp aging/ or older adulthood/ or geriatrics/ or nursing home residents/ or middle adulthood/ (98247)
- 2 (ageing or aging or care home resident* or community-dwelling or elder* or frail* or geriatric* or "late life" or "later life" or "mature adult*" or "middle age*" or "mid* life" or midlife or nonagenarian* or nursing home resident* or octogenarian* or old age* or "oldest old" or pensioner* or postmenopaus* or post-menopaus* or retired or retiree* or senior citizen* or seniors or (older adj3 adult*) or old age* or (older adj3 client*) or (older adj3 communit*) or (older adj3 female*) or (older adj3 individual*) or (older adj3 inpatient*) or (older adj3 in-patient*) or (older adj3 male*) or (older adj3 men) or (older adj3 outpatient*) or (older adj3 out-patient*) or (older adj3 patient*) or (older adj3 people) or (older adj3 person*) or (older adj3 population*) or (older adj3 women)).tw. (235261)
- 3 or/1-2 (248253)
- 4 exp communication/ or loneliness/ or social isolation/ or social support/ or friendship/ or social interaction/ (393035)
- 5 (befriend* or connectedness or (connect* adj2 other*) or (connect* adj2 people) or friendship* or interpersonal or loneliness or lonely or make friend* or making friend* or psychosocial or wellbeing or well-being).tw. (306929)
- 6 ((enhanc* or improv* or increas* or maintain*) adj5 (communicat* or connect* or interaction*)).tw. (36081)
- 7 ((social* adj2 contact*) or (social* adj2 connect*) or (social* adj2 distanc*) or (social* adj2 exclu*) or (social* adj2 inclu*) or (social* adj2 interact*) or (social* adj2 isolat*) or (social* adj2 participat*) or (social* adj2 relation*)).tw. (113124)
- 8 or/4-7 (740119)

- 9 technology.hw. (53981)
- 10 exp *computers/ or exp digital technology/ or exp electronic communication/ or exp human computer interaction/ or wireless technologies/ (157495)
- 11 (app or apps or audio* or chatbot* or chat forum* or chat interface* or chatroom* or chat room* or chat site* or chat software or computer* or digital* or DVD* or email* or e-mail* or electronic or facebook* or facetime or helpline* or help-line* or instant messag* or live chat* or messaging or mms or mobile app* or online* or phone* or phoning or robot* or short messag* service* or skype* or smartphone* or sms or social network* or social media or support line* or technolog* or telephon* or texting or text messag* or tweets or twitter or video conferenc* or videoconferenc* or video record* or videorecord* or virtual* or web app* or web-based or webcam* or webcast* or web portal* or website* or web site* or "web 2.0" or "web 3.0" or whatsapp or wireless or world wide web).ti. (133134)
- 12 (app or apps or audio* or chatbot* or chat forum* or chat interface* or chatroom* or chat room* or chat site* or chat software or computer* or digital* or DVD* or email* or e-mail* or electronic or facebook* or facetime or helpline* or help-line* or instant messag* or live chat* or messaging or mms or mobile app* or online* or phone* or phoning or robot* or short messag* service* or skype* or smartphone* or sms or social network* or social media or support line* or technolog* or telephon* or texting or text messag* or tweets or twitter or video conferenc* or videoconferenc* or video record* or videorecord* or virtual* or web app* or web-based or webcam* or webcast* or web portal* or website* or web site* or "web 2.0" or "web 3.0" or whatsapp or wireless or world wide web).ab. /freq=2 (193359)
- 13 (e-counsel* or ehealth* or e-health or e-intervention* or esupport* or e-support* or etherap* or e-therap* or mhealth* or m-health or mobile health* or online CBT or remote care or remote consult* or remote* deliver* or remote health* or self-care or self-guided or self-help or telecare or tele-care or teleconsult* or tele-consult* or telehealth* or tele-health* or telegroup* or tele-group* or telemed* or tele-med* or teletherap* or tele-therap* or virtual realit*).tw. (35201)
- 14 or/9-13 (324427)
- 15 (benefit* or change or changes or contribut* or decreas* or develop* or effect or effects or effectiveness or enhance* or evaluat* or experience* or experiment* or impact* or implement* or increas* or intervention* or method* or outcome* or pilot* or program*

- or provid* or reduc* or study or support* or system* or target* or technolog* or training
or trial or "use of ").ti. (1699614)
- 16 "systematic review"/ (602)
- 17 systematic review.md,tw. (38916)
- 18 meta analysis/ (5007)
- 19 (meta-analys* or metaanalys*).md,tw. (41526)
- 20 (cochrane or embase or medline or pubmed).ab. (27492)
- 21 exp randomized controlled trials/ (948)
- 22 randomi*.tw. (93517)
- 23 placebo.ab. (40571)
- 24 ((doubl* or singl*) adj blind*).tw. (26918)
- 25 (randomly adj2 (allocated or assigned)).ab. (42958)
- 26 trial.ti. (32761)
- 27 (group or groups).ab. (896743)
- 28 ((quasi experiment* or quasiexperiment* or quasi randomi* or quasirandomi*) adj2
(design* or method* or study or trial)).ab. (8935)
- 29 ((before adj5 after) or (controlled adj3 study) or (controlled adj3 trial) or control group* or
effect? or evaluat* or experience* or impact? or intervention* or (pre adj5 post) or
((pretest or pre-test) and (posttest or post test))).ab. /freq=2 (1158826)
- 30 (controlled before* adj2 after).tw. (120)
- 31 time series/ (2321)
- 32 (time series adj5 (analys* or design* or interrupted or ITS or studies or study or trial)).tw.
(4361)
- 33 or/16-32 (1828308)
- 34 3 and 8 and 14 and 15 and 33 (1214)
-

Database: CINAHL via EBSCO

Search Date: 16 May 2021

Yield: 1704

- S33 S32 Limiters - Exclude MEDLINE records 1,704
- S32 S3 AND S8 AND S13 AND S14 AND S31 3,725
- S31 (S29 NOT S30) 2,424,463

- S30 ((MH (animal studies) OR MH animals+ OR TI (animal model* OR mouse model* OR murine model* OR rat model*)) NOT MH (human)) 194,541
- S29 (S15 OR S16 OR S17 OR S18 OR S19 OR S20 OR S21 OR S22 OR S23 OR S24 OR S25 OR S26 OR S27 OR S28) 2,533,612
- S28 TI ((time series N5 (analys* OR design* OR interrupted OR ITS OR studies OR study OR trial))) OR AB ((time series N5 (analys* OR design* OR interrupted OR ITS OR studies OR study OR trial))) 4,680
- S27 MH ("Interrupted Time Series Analysis") 575
- S26 TI (controlled before* N2 after) OR AB (controlled before* N2 after) 852
- S25 MH ("Controlled Before-After Studies") 198
- S24 AB ((before N5 after) OR (controlled N3 study) OR (controlled N3 trial) OR control group* OR effect? OR evaluat* OR experience* OR impact* OR intervention* OR (pre N5 post) OR ((pretest OR pre-test) and (posttest OR post test))) 1,911,746
- S23 TI (((quasi experimental OR quasiexperimental OR quasi randomi* OR quasirandomi*) N2 (design* OR method* OR study OR trial))) OR AB (((quasi experimental OR quasiexperimental OR quasi randomi* OR quasirandomi*) N2 (design* OR method* OR study OR trial))) 12,524
- S22 TI (((singl* N1 blind*) OR (doubl* N1 blind*))) OR AB (((singl* N1 blind*) OR (doubl* N1 blind*))) 50,223
- S21 TI (intervention* OR trial) 207,899
- S20 AB (allocat* OR assign* OR control* OR random*) 823,932
- S19 (MH "Randomized Controlled Trials") OR (MH "Single-Blind Studies") OR (MH "Double-Blind Studies") OR (MH "Triple-Blind Studies") OR (MH "Random Assignment") OR MH ("Intervention Trials") OR MH ("Crossover Design") OR MH ("Cluster Sample") OR MH ("Placebos") 204,576
- S18 PT randomized controlled trial 128,144
- S17 AB (cochrane OR embase OR medline OR pubmed OR searched OR searches OR group OR groups) 895,315
- S16 TI ((systematic N2 review) OR TI meta analy* OR TI metaanaly*) OR AB ((systematic N2 review) OR TI meta analy* OR TI metaanaly*) 135,673
- S15 (MH "Systematic Review") OR (MH "Meta Analysis") 119,987
- S14 TI (benefit* OR change OR changes OR contribut* OR decreas* OR develop* OR effect OR effects OR effectiveness OR enhance* OR evaluat* OR experience* OR experiment* OR impact* OR implement* OR increas* OR intervention* OR method* OR outcome*

- OR pilot* OR program* OR provid* OR reduc* OR study OR support* OR system* OR target* OR technolog* OR training OR trial OR "use of ") 2,237,094
- S13 (S9 OR S10 OR S11 OR S12) 269,987
- S12 TI ((e-counsel* OR ehealth* OR e-health OR e-intervention* OR esupport* OR e-support* OR etherap* OR e-therap* OR mhealth* OR m-health OR mobile health* OR online CBT OR remote care OR remote consult* OR remote* deliver* OR remote health* OR self-care OR self-guided OR self-help OR telecare OR tele-care OR teleconsult* OR tele-consult* OR telehealth* OR tele-health* OR telegroup* OR tele-group* OR telemed* OR tele-med* OR teletherap* OR tele-therap* OR virtual realit*)) OR AB ((e-counsel* OR ehealth* OR e-health OR e-intervention* OR esupport* OR e-support* OR etherap* OR e-therap* OR mhealth* OR m-health OR mobile health* OR online CBT OR remote care OR remote consult* OR remote* deliver* OR remote health* OR self-care OR self-guided OR self-help OR telecare OR tele-care OR teleconsult* OR tele-consult* OR telehealth* OR tele-health* OR telegroup* OR tele-group* OR telemed* OR tele-med* OR teletherap* OR tele-therap* OR virtual realit*))47,907
- S11 TI ((app OR apps OR audio* OR chatbot* OR chat forum* OR chat interface* OR chatroom* OR chat room* OR chat site* OR chat software OR computer* OR digital* OR DVD* OR email* OR e-mail* OR electronic OR facebook* OR facetime OR helpline* OR help-line* OR instant messag* OR live chat* OR messaging OR mms OR mobile app* OR online* OR phone* OR phoning OR robot* OR short messag* service* OR skype* OR smartphone* OR sms OR social network* OR social media OR support line* OR technolog* OR telephon* OR texting OR text messag* OR tweets OR twitter OR video conferenc* OR videoconferenc* OR video record* OR videorecord* OR virtual* OR web app* OR web-based OR webcam* OR webcast* OR web portal* OR website* OR web site* OR "web 2.0" OR "web 3.0" OR whatsapp OR wireless OR world wide web)) AND AB ((app OR apps OR audio* OR chatbot* OR chat forum* OR chat interface* OR chatroom* OR chat room* OR chat site* OR chat software OR computer* OR digital* OR DVD* OR email* OR e-mail* OR electronic OR facebook* OR facetime OR helpline* OR help-line* OR instant messag* OR live chat* OR messaging OR mms OR mobile app* OR online* OR phone* OR phoning OR robot* OR short messag* service* OR skype* OR smartphone* OR sms OR social network* OR social media OR support line* OR technolog* OR telephon* OR texting OR text messag* OR tweets OR twitter OR video conferenc* OR videoconferenc* OR video record* OR videorecord* OR virtual* OR web

- app* OR web-based OR webcam* OR webcast* OR web portal* OR website* OR web site* OR "web 2.0" OR "web 3.0" OR whatsapp OR wireless OR world wide web))89,765
- S10 (MM "Computers and Computerization") OR (MM "Computer Environment+") OR (MM "Computer Systems+") OR (MM "Mobile Applications") OR (MM "Telecommunications+") OR (MH "Therapy, Computer Assisted") 120,310
- S9 MW technology 73,772
- S8 (S4 OR S5 OR S6 OR S7) 367,212
- S7 TI (((social* N2 contact*) OR (social* N2 connect*) OR (social* N2 distanc*) OR (social* N2 exclu*) OR (social* N2 inclu*) OR (social* N2 interact*) OR (social* N2 isolat*) OR (social* N2 participat*) OR (social* N2 relation*))) OR AB (((social* N2 contact*) OR (social* N2 connect*) OR (social* N2 distanc*) OR (social* N2 exclu*) OR (social* N2 inclu*) OR (social* N2 interact*) OR (social* N2 isolat*) OR (social* N2 participat*) OR (social* N2 relation*))) 42,336
- S6 TI (((enhanc* OR improv* OR increas* OR maintain*) N5 (communicat* OR connect* OR interaction*))) OR AB (((enhanc* OR improv* OR increas* OR maintain*) N5 (communicat* OR connect* OR interaction*))) 28,433
- S5 TI ((befriend* OR connectedness OR (connect* N2 other*) OR (connect* N2 people) OR friendship* OR interpersonal OR loneliness OR lonely OR make friend* OR making friend* OR psychosocial OR wellbeing OR well-being)) OR AB ((befriend* OR connectedness OR (connect* N2 other*) OR (connect* N2 people) OR friendship* OR interpersonal OR loneliness OR lonely OR make friend* OR making friend* OR psychosocial OR wellbeing OR well-being)) 142,617
- S4 (MH Communication) OR (MH "Social Isolation+") OR (MH "Support, Psychosocial") OR (MH "Interpersonal Relations") OR (MH "Friendship") OR (MH "Social Integration") 215,242
- S3 (S1 OR S2) 1,448,300
- S2 TI ((ageing OR aging OR care home resident* OR centenarian* OR community-dwelling OR elder* OR frail* OR geriatric* OR "late life" OR "later life" OR "mature adult*" OR "middle age*" OR "mid* life" OR midlife OR nonagenarian* OR nursing home resident* OR octogenarian* OR old age* OR "oldest old" OR pensioner* OR postmenopaus* OR post-menopaus* OR retired OR retiree* OR senior* OR septuagenarian OR sexagenarian* OR veteran*)) OR AB ((ageing OR aging OR care home resident* OR centenarian* OR community-dwelling OR elder* OR frail* OR geriatric* OR "late life" OR "later life" OR "mature adult*" OR "middle age*" OR "mid* life" OR midlife OR

nonagenarian* OR nursing home resident* OR octogenarian* OR old age* OR "oldest old" OR pensioner* OR postmenopaus* OR post-menopaus* OR retired OR retiree* OR senior* OR septuagenarian OR sexagenarian* OR veteran*)) 300,074

S1 (MH "Aged+") OR (MH "Middle Age") 1,326,993

Database: Web of Science via Clarivate (Indexes=SCI-EXPANDED, SSCI, CPCI-S, CPCI-SSH, ESCI) Timespan=All years

Search Date: 17 May 2021

Yield: 853

#8 853 #7 AND #6 AND #5 AND #4

#7 2,072,148 TS=(controlled before-after OR "controlled trial" OR "doubl* blind*" OR "meta-analy*" OR "metaanaly*" OR randomi* OR randomly OR "singl* blind*" OR "systematic review" OR "time series")

#6 1,917,703 TI=(app OR apps OR audio* OR chatbot* OR "chat forum*" OR "chat interface*" OR chatroom* OR "chat room*" OR "chat site*" OR "chat software" OR computer* OR digital* OR DVD* OR "e-counsel*" OR ehealth* OR "e-health" OR "e-intervention*" OR email* OR e-mail* OR electronic OR esupport* OR "e-support*" OR etherap* OR "e-therap*" OR facebook* OR facetime OR helpline* OR "help line*" OR "instant messag*" OR "live chat*" OR mhealth* OR "m-health" OR "mobile health*" OR messaging OR mms OR "mobile app*" OR online* OR phone* OR phoning OR "remote care" OR "remote consult*" OR "remote* deliver*" OR "remote health*" OR robot* OR "self-care" OR "self-guided" OR "self-help" OR "short messag* service*" OR skype* OR smartphone* OR sms OR "social network*" OR "social media" OR "support line*" OR technolog* OR telecare OR "tele-care" OR teleconsult* OR "tele-consult*" OR telehealth* OR "tele-health*" OR telegroup* OR "tele-group*" OR telemed* OR "tele-med*" OR telephon* OR teletherap* OR "tele-therap*" OR texting OR "text messag" OR tweets OR twitter OR "video conferenc*" OR videoconferenc* OR virtual* OR "web app*" OR "web-based" OR webcam* OR webcast* OR "web portal*" OR website* OR "web site*" OR "web 2.0" OR "web 3.0" OR whatsapp OR wireless OR "world wide web")

#5 762,426 TI=(befriend* OR connected* OR loneliness OR lonely OR psychosocial* OR "social isolation" OR "social support*" OR "socially isolated" OR "well-being" OR wellbeing) OR AB=(befriend* OR connected* OR loneliness OR lonely OR psychosocial*

OR "social isolation" OR "social support*" OR "socially isolated" OR "well-being" OR wellbeing)

#4 4,257,374 #3 OR #2 OR #1

#3 1,062,970 TI=((older NEAR/3 adult*) OR (older NEAR/3 client*) OR (older NEAR/3 communit*) OR (older NEAR/3 female*) OR (older NEAR/3 individual*) OR (older NEAR/3 inpatient*) OR (older NEAR/3 in-patient*) OR (older NEAR/3 male*) OR (older NEAR/3 men) OR (older NEAR/3 outpatient*) OR (older NEAR/3 out-patient*) OR (older NEAR/3 patient*) OR (older NEAR/3 people) OR (older NEAR/3 person*) OR (older NEAR/3 population*) OR (older NEAR/3 women)) OR AB=((older NEAR/3 adult*) OR (older NEAR/3 client*) OR (older NEAR/3 communit*) OR (older NEAR/3 female*) OR (older NEAR/3 individual*) OR (older NEAR/3 inpatient*) OR (older NEAR/3 in-patient*) OR (older NEAR/3 male*) OR (older NEAR/3 men) OR (older NEAR/3 outpatient*) OR (older NEAR/3 out-patient*) OR (older NEAR/3 patient*) OR (older NEAR/3 people) OR (older NEAR/3 person*) OR (older NEAR/3 population*) OR (older NEAR/3 women) OR (old age))

#2 302.219 SU=(aged OR ageing OR aging OR elder* OR geriatric* OR middle age OR older OR senior*)

#1 3,686,745 TI=(ageing OR aging OR "care home resident*" OR centenarian* OR "community-dwelling" OR elder* OR frail* OR geriatric* OR "late life" OR "later life" OR "mature adult*" OR "middle age*" OR "mid* life" OR midlife OR nonagenarian* OR "nursing home resident*" OR octogenarian* OR "old age*" OR "oldest old" OR pensioner* OR postmenopaus* OR post-menopaus* OR retired OR retiree* OR senior* OR septuagenarian OR sexagenarian* OR veteran*) OR AB=(aged OR ageing OR aging OR "care home resident*" OR centenarian* OR "community-dwelling" OR elder* OR frail* OR geriatric* OR "late life" OR "later life" OR "mature adult*" OR "middle age*" OR "mid* life" OR midlife OR nonagenarian* OR "nursing home resident*" OR octogenarian* OR "old age*" OR "oldest old" OR pensioner* OR postmenopaus* OR post-menopaus* OR retired OR retiree* OR senior* OR septuagenarian OR sexagenarian* OR veteran*)

Database: ProQuest (all databases)

Search Date: 17 May 2021

Yield: 706

ti(("care home resident*" OR "community-dwelling" OR elder* OR frail OR geriatric* OR "late life" OR "later life" OR "middle-age*" OR mid-life OR midlife OR "nursing home resident*" OR "old age*" OR older OR pensioner* OR retired OR retiree* OR "senior citizen*" OR seniors)) AND (controlled before-after OR controlled trial OR "meta-analysis" OR randomi* OR randomly OR "systematic review" OR "time series") AND ti((befriend* OR connected* OR loneliness OR lonely OR psychosocial* OR "social isolation" OR "social support*" OR "socially isolated" OR "well-being" OR wellbeing)) AND noft((app OR apps OR audio* OR chatbot* OR chat forum* OR chat interface* OR chatroom* OR "chat room*" OR chat site* OR chat software OR computer* OR digital* OR DVD* OR email* OR e-mail* OR electronic OR facebook* OR facetime OR helpline* OR help-line* OR instant messag* OR live chat* OR messaging OR mms OR mobile app* OR online* OR phone* OR phoning OR robot* OR short messag* service* OR skype* OR smartphone* OR sms OR social network* OR social media OR support line* OR technolog* OR telephon* OR texting OR text messag* OR tweets OR twitter OR video conferenc* OR videoconferenc* OR video record* OR videorecord* OR virtual* OR web app* OR web-based OR webcam* OR webcast* OR "web portal*" OR website* OR "web site*" OR "web 2.0" OR "web 3.0" OR whatsapp OR wireless OR "world wide web"))

Database: International Bibliography of the Social Sciences (IBSS) via ProQuest

Search Date: 17 May 2021

Yield: 275

noft(("care home resident*" OR "community-dwelling" OR elder* OR frail OR geriatric* OR "late life" OR "later life" OR "middle-age*" OR mid-life OR midlife OR "nursing home resident*" OR "old age*" OR older OR pensioner* OR retired OR retiree* OR "senior citizen*" OR seniors)) AND noft((app OR apps OR audio* OR chatbot* OR chat forum* OR chat interface* OR chatroom* OR "chat room*" OR chat site* OR chat software OR computer* OR digital* OR DVD* OR email* OR e-mail* OR electronic OR facebook* OR facetime OR helpline* OR help-line* OR instant messag* OR live chat* OR messaging OR mms OR mobile app* OR online* OR phone* OR phoning OR robot* OR short messag* service* OR skype* OR smartphone* OR sms OR social network* OR social media OR support line* OR technolog* OR telephon* OR texting OR text messag* OR

tweets OR twitter OR video conferenc* OR videoconferenc* OR video record* OR
videorecord* OR virtual* OR web app* OR web-based OR webcam* OR webcast* OR
"web portal*" OR website* OR "web site*" OR "web 2.0" OR "web 3.0" OR whatsapp OR
wireless OR "world wide web")) AND noft((befriend* OR connected* OR loneliness OR
lonely OR psychosocial* OR "social isolation" OR "social support*" OR "socially isolated"
OR "well-being" OR wellbeing)) AND (controlled before-after OR controlled trial OR
"meta-analysis" OR randomi* OR randomly OR "systematic review" OR "time series")

Database: EBSCO (all databases except CINAHL)

Search Date: 17 May 2021

Yield: 578

TX (("care home resident*" OR "community-dwelling" OR elder* OR frail OR geriatric* OR
"late life" OR "later life" OR "middle-age*" OR mid-life OR midlife OR "nursing home
resident*" OR "old age*" OR older OR pensioner* OR retired OR retiree* OR "senior
citizen*" OR seniors)) AND TI ((befriend* OR connected* OR loneliness OR lonely OR
psychosocial* OR "social isolation" OR "social support*" OR "socially isolated" OR "well-
being" OR wellbeing)) AND TI ((app OR apps OR audio* OR chatbot* OR chat forum*
OR chat interface* OR chatroom* OR "chat room*" OR chat site* OR chat software OR
computer* OR digital* OR DVD* OR email* OR e-mail* OR electronic OR facebook* OR
facetime OR helpline* OR help-line* OR instant messag* OR live chat* OR messaging
OR mms OR mobile app* OR online* OR phone* OR phoning OR robot* OR short
messag* service* OR skype* OR smartphone* OR sms OR social network* OR social
media OR support line* OR technolog* OR telephon* OR texting OR text messag* OR
tweets OR twitter OR video conferenc* OR videoconferenc* OR video record* OR
videorecord* OR virtual* OR web app* OR web-based OR webcam* OR webcast* OR
"web portal*" OR website* OR "web site*" OR "web 2.0" OR "web 3.0" OR whatsapp OR
wireless OR "world wide web")) AND TX ((controlled before-after OR controlled trial
OR "meta-analysis" OR randomi* OR randomly OR "systematic review" OR "time
series"))

Database: Global Index Medicus

Search Date: 17 May 2021

Yield: 213

tw:((tw:(("care home resident*" OR "community-dwelling" OR elder* OR frail OR geriatric* OR "late life" OR "later life" OR "middle-age*" OR mid-life OR midlife OR "nursing home resident*" OR "old age*" OR older OR pensioner* OR retired OR retiree* OR "senior citizen*" OR seniors))) AND (tw:(befriend* OR connected* OR loneliness OR lonely OR psychosocial* OR "social isolation" OR "social support*" OR "socially isolated" OR "well-being" OR wellbeing))) AND (ti:(app OR apps OR audio* OR chatbot* OR "chat forum*" OR "chat interface*" OR chatroom* OR "chat room*" OR "chat site*" OR "chat software" OR computer* OR digital* OR dvd* OR "e-counsel*" OR ehealth* OR "e-health" OR "e-intervention*" OR email* OR e-mail* OR electronic OR esupport* OR "e-support*" OR etherap* OR "e-therap*" OR facebook* OR facetime OR helpline* OR "help line*" OR "instant messag*" OR "live chat*" OR mhealth* OR "m-health" OR "mobile health*" OR messaging OR mms OR "mobile app*" OR online* OR phone* OR phoning OR "remote care" OR "remote consult*" OR "remote* deliver*" OR "remote health*" OR robot* OR "self-care" OR "self-guided" OR "self-help" OR "short messag* service*" OR skype* OR smartphone* OR sms OR "social network*" OR "social media" OR "support line*" OR technolog* OR telecare OR "tele-care" OR teleconsult* OR "tele-consult*" OR telehealth* OR "tele-health*" OR telegroup* OR "tele-group*" OR telemed* OR "tele-med*" OR telephon* OR teletherap* OR "tele-therap*" OR texting OR "text messag" OR tweets OR twitter OR "video conferenc*" OR videoconferenc* OR virtual* OR "web app*" OR "web-based" OR webcam* OR webcast* OR "web portal*" OR website* OR "web site*" OR "web 2.0" OR "web 3.0" OR whatsapp OR wireless OR "world wide web"))))

Database: Epistemonikos

Search Date: 17 May 2021

Yield: 1093

(title:(ageing OR aging OR "care home resident*" OR "community-dwelling" OR elder* OR frail OR geriatric* OR "late life" OR "later life" OR "middle-age*" OR mid-life OR midlife OR

"nursing home resident*" OR "old age*" OR older OR pensioner* OR retired OR retiree* OR "senior citizen*" OR seniors)) OR abstract:((ageing OR aging OR "care home resident*" OR "community-dwelling" OR elder* OR frail OR geriatric* OR "late life" OR "later life" OR "middle-age*" OR mid-life OR midlife OR "nursing home resident*" OR "old age*" OR older OR pensioner* OR retired OR retiree* OR "senior citizen*" OR seniors))) AND (title:((befriend* OR connected* OR loneliness OR lonely OR psychosocial* OR "social isolation" OR "social support*" OR "socially isolated" OR "well-being" OR wellbeing)) OR abstract:((befriend* OR connected* OR loneliness OR lonely OR psychosocial* OR "social isolation" OR "social support*" OR "socially isolated" OR "well-being" OR wellbeing))) AND (title:((app OR apps OR audio* OR chatbot* OR "chat forum*" OR "chat interface*" OR chatroom* OR "chat room*" OR "chat site*" OR "chat software" OR computer* OR digital* OR DVD* OR "e-counsel*" OR ehealth* OR "e-health" OR "e-intervention*" OR email* OR e-mail* OR electronic OR esupport* OR "e-support*" OR etherap* OR "e-therap*" OR facebook* OR facetime OR helpline* OR "help line*" OR "instant messag*" OR "live chat*" OR mhealth* OR "m-health" OR "mobile health*" OR messaging OR mms OR "mobile app*" OR online* OR phone* OR phoning OR "remote care" OR "remote consult*" OR "remote* deliver*" OR "remote health*" OR robot* OR "self-care" OR "self-guided" OR "self-help" OR "short messag* service*" OR skype* OR smartphone* OR sms OR "social network*" OR "social media" OR "support line*" OR technolog* OR telecare OR "tele-care" OR teleconsult* OR "teleconsult*" OR telehealth* OR "tele-health*" OR telegroup* OR "tele-group*" OR telemed* OR "tele-med*" OR telephon* OR teletherap* OR "tele-therap*" OR texting OR "text messag" OR tweets OR twitter OR "video conferenc*" OR videoconferenc* OR virtual* OR "web app*" OR "web-based" OR webcam* OR webcast* OR "web portal*" OR website* OR "web site*" OR "web 2.0" OR "web 3.0" OR whatsapp OR wireless OR "world wide web")) OR abstract:((app OR apps OR audio* OR chatbot* OR "chat forum*" OR "chat interface*" OR chatroom* OR "chat room*" OR "chat site*" OR "chat software" OR computer* OR digital* OR DVD* OR "e-counsel*" OR ehealth* OR "e-health" OR "e-intervention*" OR email* OR e-mail* OR electronic OR esupport* OR "e-support*" OR etherap* OR "e-therap*" OR facebook* OR facetime OR helpline* OR "help line*" OR "instant messag*" OR "live chat*" OR mhealth* OR "m-health" OR "mobile health*" OR messaging OR mms OR "mobile app*" OR online* OR phone* OR phoning OR "remote care" OR "remote consult*" OR "remote* deliver*" OR "remote health*" OR robot* OR "self-care" OR "self-guided" OR "self-help" OR "short messag*

service*" OR skype* OR smartphone* OR sms OR "social network*" OR "social media" OR "support line*" OR technolog* OR telecare OR "tele-care" OR teleconsult* OR "tele-consult*" OR telehealth* OR "tele-health*" OR telegroup* OR "tele-group*" OR teledem* OR "tele-med*" OR telephon* OR teletherap* OR "tele-therap*" OR texting OR "text messag" OR tweets OR twitter OR "video conferenc*" OR videoconferenc* OR virtual* OR "web app*" OR "web-based" OR webcam* OR webcast* OR "web portal*" OR website* OR "web site*" OR "web 2.0" OR "web 3.0" OR whatsapp OR wireless OR "world wide web"))

Publication type: Systematic Review OR Primary Study

3. Detailed eligibility criteria

Criteria	Inclusion	Exclusion
Population	<ul style="list-style-type: none"> - Older adults ≥ 60 years old - Mixed population with disaggregated data for target population (≥ 60 years old) - Participants in community settings (personal homes, residential homes, or independent living facilities) or supportive care institutions (nursing homes or long-term care and assisted living facilities). - *Systematic reviews that are focused on older adults without specifying age 	<ul style="list-style-type: none"> - Not older adults (< 60 years old) - Mixed population without disaggregated data for target population (≥ 60 years old) - Hospitalized patients
Intervention	<p>Digital interventions focusing on SIL:</p> <ul style="list-style-type: none"> - Social skills training (e.g., computer and internet training) - Enhancing social support (e.g., telephone befriending, telecare, robopets) - Increasing social interaction (e.g., videoconferencing, social networking sites, chat box) - Social cognitive training (e.g., online CBT with a focus on reducing SIL) <p>If a systematic review has a more general focus and looks at all intervention types, not</p>	<ul style="list-style-type: none"> - digital interventions with no focus on SIL e.g. e-health exclusively for healthcare - digital interventions for monitoring (e.g., sensors for falls) and smart homes - interventions taking place in hospital settings

	just digital interventions, it will be included if the data is disaggregated by intervention type (shows results for digital interventions separately).	
Evidence/objective	Focus on SIL, mental health and well-being	Not focused on SIL (e.g., CBT for dementia care)
Study design	<ul style="list-style-type: none"> - Systematic reviews - Randomized controlled trials - Quasi-experimental studies - Controlled before-after studies - Interrupted time series with at least six data points (3 before and 3 after a discrete intervention) - Regression discontinuity designs - *Studies with a comparison/control group - Protocols of systematic reviews and eligible primary studies 	<ul style="list-style-type: none"> - Not a systematic review (e.g., literature reviews) – does not satisfy at least 4/5 criteria (i) Were inclusion/exclusion criteria reported? (ii) Was the search adequate? (iii) Were the included studies synthesized? (iv) Was the quality of the included studies assessed; (v) Are sufficient details about the individual included studies presented? - less than six period interrupted time series design - longitudinal cohort studies with no controls - Cross-sectional studies - Qualitative study designs

4. EGM Coding tool template

Category	Subcategories
Study characteristics	
Publication status	<ul style="list-style-type: none"> • Complete • On-going (e.g., protocols, trial registrations) • Conference abstract
Study design	<ul style="list-style-type: none"> • Primary study • Systematic review
Primary study design	<ul style="list-style-type: none"> • Randomized controlled trials (RCTs) • non-randomized studies
Equity focus: Is the study population identified by the authors as aimed at/focused	<ul style="list-style-type: none"> • Place of residence • Race/Ethnicity

<p>on disadvantaged across any PROGRESS factors</p>	<ul style="list-style-type: none"> • Occupation • Gender/sex • Religion • Education • Socioeconomic status • Social capital (e.g. living alone) • Plus factor - Frailty • Plus factor - Digital literacy • Plus factor - Health status e.g. dementia, disease severity • Plus factor - Socially isolated or at risk of social isolation • Plus factor - Lonely or at risk of loneliness • Plus factor - Age • Plus factor - Disability Plus factor - Living situation (e.g. alone, long-term care)
<p>If population is identified as “at risk”, how are they identified</p>	<ul style="list-style-type: none"> • Case-finding • Outreach • From a community-based program • Screening in primary care
<p>Methodological quality assessment of systematic reviews</p>	<ul style="list-style-type: none"> • Low or critically low quality reviews • Moderate quality reviews • High quality reviews
<p>Interventions</p>	
<p>Improving social skills</p>	<ul style="list-style-type: none"> • Training in how to use digital technology (e.g., computer, internet) • Digitally delivered training (e.g. about caregiving/skills building) • Digitally delivered learning (e.g. new language)
<p>Enhancing social interaction</p>	<ul style="list-style-type: none"> • Digital connections with family/friends • Digital connections with community
<p>Enhancing social support</p>	<ul style="list-style-type: none"> • Digital/remote e-health services

	<ul style="list-style-type: none"> • Digital social and health care coordination with family/friends • Geolocating/identifying older adults who need services • Socially assistive robots and virtual pets • Virtual spaces • Virtual assistants (e.g., Google home, Alexa) • Virtual social support groups • Digital intergenerational approaches • Digital games (e.g., scrabble, chess, cards, exergames) • Digitally delivered activities (e.g., exercise - yoga, tai chi) • Digital coordination of health or social care services (e.g., online referrals)
Social cognitive training	<ul style="list-style-type: none"> • Digital cognitive behavioural therapy • Digital mindfulness training • Digital psychoeducation • Digital reminiscence therapy • Digital cognitive behavioural coaching
Multicomponent	
Intervention focus	<ul style="list-style-type: none"> • Loneliness • Social isolation • Social isolation and loneliness
Intervention format	<ul style="list-style-type: none"> • Group-based • One-on-one
How technologies are used	<ul style="list-style-type: none"> • Telephone calls • E-mail • Messaging • Discussion forums • Videoconferencing

	<ul style="list-style-type: none"> • Social networking sites e.g., Facebook, Twitter • Digital games • Robots and virtual companions • Virtual assistants • Virtual spaces or classrooms with messaging capabilities • Mobility tools (e.g. walky - microblogging walking frame) • Participating in an activity • Music • Unspecified
How training on how to use of digital technology was delivered	<ul style="list-style-type: none"> • Digitally (remotely) delivered training • In-person (face to face) training • Unspecified
Outcomes	
Individual	<ul style="list-style-type: none"> • Loneliness • Social isolation • Social connectedness • Quality of life/well-being • Anxiety/depression • Confidence level or self esteem • Adverse effects
Community	<ul style="list-style-type: none"> • Social support • Social engagement • Social cohesion • Social capital • Digital divide
Process indicators	<ul style="list-style-type: none"> • Acceptance • Adherence • Technology use • Feasibility

	<ul style="list-style-type: none"> • Cost-effectiveness) • Affordability • Barriers
Population sociodemographics	
Age groups	<ul style="list-style-type: none"> • includes 60-70 years • includes 70-80 years • includes >80 years • Restricted to >80 years • Unspecified
Health conditions	<ul style="list-style-type: none"> • Communicable disease • Noncommunicable disease • Dementia/Alzheimer's disease • Depression • Comorbidity • Disability • Physical frailty • Psychological frailty • Social frailty • Care dependent • Discharge from hospital • End-of-life/palliative care
PROGRESS-Plus factors	<ul style="list-style-type: none"> • Place of residence (urban/rural or remote/unspecified) • Race/Ethnicity • Occupation • Gender or Sex (men only/women only/LGBTQIA2S+/unspecified) • Religion • Education • Socioeconomic status • Social capital (marital status) • Living alone

	<ul style="list-style-type: none"> • Caregivers •
Needs	<ul style="list-style-type: none"> • Social and emotional needs • Purpose in life • Mobility • Personal care needs • Meals • Domestic assistance • Accommodation (housing/home modifications and maintenance) • Financial management • Communication (language support/interpreters, information and assistance/referral services) • Skills development • Learning (e.g., a new activity, language or about social skills) • Care navigation support or task orientation • Clinical/health needs • Respite care • Caregiver support
Setting	<ul style="list-style-type: none"> • Personal home • Independent living (residential home) • Assisted living • Long-term care/nursing home • Unspecified
World Health Organization regions	<ul style="list-style-type: none"> • African Region • Regions of the Americas • South-East Asian Region • European Region • Eastern Mediterranean Region • Western Pacific Region

	<ul style="list-style-type: none"> • Multiple regions • Unspecified
World Bank Classification by income	<ul style="list-style-type: none"> • Low-income economies • Lower-middle income economies • Upper-middle income economies • High-income economies • Unspecified
Countries	<ul style="list-style-type: none"> • Africa (where multiple countries) • Europe (where multiple countries) • Afghanistan • Angola • Armenia • Argentina • Australia • Austria • Bahamas, The • Bahrain • Bangladesh • Barbados • Belgium • Belarus • Belize • Bolivia • Brazil • Botswana • Bulgaria • Burkina Faso • Cambodia • Cameroon • Canada • Chile • China

- Congo
- Croatia
- Cuba
- Colombia
- Cyprus
- Czech Republic
- Dominican Republic
- Denmark
- Egypt
- Ecuador
- Eritrea
- Estonia
- Finland
- Ethiopia
- Gambia, The
- France
- Georgia
- Ghana
- Germany
- Greece
- Guatemala
- Guinea-Bissau
- Haiti
- Honduras
- Hong Kong
- Hungary
- Iceland
- India
- Indonesia
- Iran
- Ireland
- Israel

- Italy
- Ivory Coast
- Kazakhstan
- Jordan
- Kenya
- Korea
- Kuwait
- Japan
- Jamaica
- Lao
- Latin America (where multiple countries)
- Lebanon
- Lesotho
- Liberia
- Latvia
- Lithuania
- Luxembourg
- Macedonia
- Malawi
- Madagascar
- Malaysia
- Mali
- Mexico
- Micronesia
- Marshall Islands
- Mozambique
- Mongolia
- Myanmar (Burma)
- Namibia
- Nepal
- New Zealand
- The Netherlands

- Nicaragua
- Niger
- Nigeria
- Northern Ireland
- Pakistan
- Norway
- Panama
- Papua New Guinea
- Peru
- Philippines
- Poland
- Portugal
- Puerto Rico
- Romania
- Russia
- Rwanda
- Samoa
- San Marino
- Saudi Arabia
- Scotland
- Serbia
- Senegal
- Singapore
- Sierra Leone
- Slovakia
- Slovenia
- South Africa
- Spain
- Sri Lanka
- St Lucia
- Swaziland
- Sweden

	<ul style="list-style-type: none"> • Switzerland • Syria • Taiwan • Thailand • Tanzania • Trinidad and Tobago • Tunisia • Uganda • Ukraine • Turkey • UK • USA • Uzbekistan • Vanuatu • Venezuela • Vietnam • West Indies • Yemen • Zambia • Zimbabwe
<p>Equity analysis: Does the study assess any differences in effects (benefit or harm) across any PROGRESS factors</p>	<ul style="list-style-type: none"> • Place of residence (rural or remote/urban) • Race/ethnicity/culture/language • Occupation • Gender/sex • Religion • Education • Socioeconomic status • Social capital (e.g. social network) • Plus factor - Frailty • Plus factor - Digital literacy • Plus factor - Health status (e.g. dementia, disease severity)

	<ul style="list-style-type: none">• Plus factor - Socially isolated or at risk• Plus factor - Lonely or at risk• Plus factor - Age• Plus factor - Disability• Plus factor - Living situation (e.g. alone, long term care)
--	---